

BIRCH, STEWART, KOLASCH & BIRCH, LLP

INTELLECTUAL PROPERTY LAW

8110 GATEHOUSE ROAD

SUITE 500 EAST

FALLS CHURCH, VA 22042-1210

USA

(703) 205-8000

FAX: (703) 205-8050

(703) 698-8590 (G IV)

e-mail: mailroom@bskb.com

web: http://www.bskb.com

CALIFORNIA OFFICE

650 TOWN CENTER DRIVE, SUITE 1120

COSTA MESA, CA 92626-7125

GARY D. YACURA
THOMAS S. AUCHTERLONIE
MICHAEL R. CAMMARATA
JAMES T. ELLER, JR.
SCOTT L. LOWE
MARY ANN CAPRIA
MARK J. NUELLE, PH.D.
DARIN E. BARTHOLOMEW*
D. RICHARD ANDERSON
PAUL C. LEWIS
W. KARL RENNER
MARK W. MILSTEAD*
JOHN CAMPA*

REG. PATENT AGENTS:
FREDERICK R. HANDREN
ANDREW J. TELESZ, JR.
MARYANNE ARMSTRONG, PH.D.
MAKI HATSUMI
MIKE S. RYU
CRAIG A. McROBBIE
GARTH M. DAHLEN, PH.D.
LAURA C. LUTZ
ROBERT E. GOOZNER, PH.D.
HYUNG N. SOHN
MATTHEW J. LATTIG
ALAN PEDERSEN-GILES
JUSTIN D. KARJALA

TERRELL C. BIRCH
RAYMOND C. STEWART
JOSEPH A. KOLASCH
JAMES M. SLATTERY
BERNARD L. SWEENEY*
MICHAEL K. MUTTER
CHARLES GORENSTEIN
GERALD M. MURPHY, JR.
LEONARD R. SVENSSON
TERRY L. CLARK
ANDREW D. MEIKLE
MARC S. WEINER
JOE MCKINNEY MUNCY
ROBERT J. KENNEY
DONALD J. DALEY
JOHN W. BAILEY
JOHN A. CASTELLANO, III
OF COUNSEL
HERBERT M. BIRCH (1905-1996)
ELLIOT A. GOLDBERG*
WILLIAM L. GATES*
EDWARD H. VALANCE
RUPERT J. BRADY (RET.)*

NOT TO BE REPRODUCED OR TRANSMITTED IN ANY FORM OR BY ANY MEANS, ELECTRONIC OR MECHANICAL, INCLUDING PHOTOCOPYING, RECORDING, OR BY ANY INFORMATION STORAGE AND RETRIEVAL SYSTEM, WITHOUT PERMISSION IN WRITING FROM THE UNITED STATES PATENT AND TRADEMARK OFFICE.

Date: December 16, 1999

Docket No.: 2121-154P

Honorable Commissioner of Patents
Washington, D.C. 20231

Sir:

This is a Request for filing a continuation X divisional
application under 37 C.F.R. § 1.53(b) of pending prior application
Serial No. 08/817,188 filed on May 15, 1997

by
Marc DeBlock

for
Genetic Transformation Using a PARP Inhibitor

- X Enclosed is a true copy of the complete prior above-identified application as filed, consisting of specification, claims and declaration.
- X The filing fee has been calculated as follows:

			LARGE ENTITY	SMALL ENTITY
BASIC FEE			\$760.00	\$380.00
	NUMBER FILED	NUMBER EXTRA	RATE FEE	RATE FEE
TOTAL CLAIMS	13- 20 =	0	x 18 = \$	x 9 = \$
INDEPENDENT CLAIMS	1- 3 =	0	x 78 = \$	x 39 = \$
MULTIPLE DEPENDENT CLAIMS PRESENTED			+ \$260.00	+ \$135.00
TOTAL			\$760.00	

3. ☒ A check in the amount of \$ 760.00 to cover the filing fee and recording fee (if applicable) is enclosed.
4. ☐ Please charge Deposit Account No. 02-2448 in the amount of \$ _____. A triplicate copy of this request is enclosed.
5. ☒ Amend the specification by inserting before the first line thereof the following:

--This application is a ____ continuation ☒ divisional of copending application Serial No. 08/817,188, filed on May 15, 1997, Application Serial No. 08/817,188 is a 371 national application of international application PCT/EP96/03366, filed on July 31, 1996. The entire contents of these applications are hereby incorporated by reference.--
6. ☐ Transfer the drawings from the prior application to this application and abandon said prior application as of the filing date accorded this application. A duplicate copy of this request is enclosed for filing in the prior application file.
7. ☐ Enclosed is/are _____ sheet(s) of _____ drawings.
8. ☐ A verified statement claiming small entity status was

filed in prior application Serial No. _____
on _____. See attached copy
of verified statement claiming small entity.

9. X The prior application is assigned to Plant Genetic
Systems, N.V.

10. ✓X A Preliminary Amendment is enclosed.

11a. ____ Priority of Application No(s). _____ filed in _____ on
_____ is/are claimed under 35 U.S.C. § 119. See
attached copy of the Letter claiming priority filed in
the prior application on _____.

11b. X Priority of International Appln. PCT/EP96/03366 filed
on July 31, 1996 under the Patent Cooperation Treaty
and European Application No. 95401844.6 filed in
Europe on August 4, 1995 under 35 U.S.C. § 119 are
hereby reclaimed.

12. ✓X An Information Disclosure Statement and PTO-1449
form(s) are attached hereto for the Examiner's
consideration.

13. X Address all future communications to:

BIRCH, STEWART, KOLASCH & BIRCH, LLP
P.O. Box 747
Falls Church, Virginia 22040-0747
Telephone: (703) 205-8000

or

Customer No. 2292

14. ____ An extension of time for _____ month(s)
until _____ has been submitted
in parent application Serial No. _____
in order to establish copendency with the present
application.

15. ____ Also enclosed herewith is the following:


Rule 53(b) Div. of 08/817,188
New Attorney Docket No.: 2121-154P

If necessary, the Commissioner is hereby authorized to charge payment of the following fees associated with this communication or credit any overpayment to Deposit Account No. 02-2448, including any additional filing fee required under 37 C.F.R. § 1.16 or any patent application processing fee under 37 C.F.R. § 1.17.

Respectfully submitted,

BIRCH, STEWART, KOLASCH & BIRCH, LLP

By



Gerald M. Murphy, Jr.
Reg. No. 28,977

P.O. Box 747
Falls Church, VA 22040-0747
(703) 205-8000

 GMM/MAA/bsh

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Marc DeBlock
Serial No: NEW-Rule 53(b) Div.
of 08/817,188 Group: Unassigned
Filed: December 16, 1999 Examiner: Unassigned
For: GENETIC TRANSFORMATION USING A PARP INHIBITOR

PRELIMINARY AMENDMENT

Assistant Commissioner of Patent
Washington, D.C. 20231

December 16, 1999

Sir:

The following preliminary amendments and remarks are respectfully submitted in connection with the above-identified application.

IN THE ABSTRACT:

Please add the attached Abstract to the end of the application now on file.

IN THE SPECIFICATION:

Please amend the specification as follows:

Please delete pages 39-60 of the specification containing the sequence listing. Please renumber the remaining pages of the specification, beginning with the claims, consecutively from page 39 of the specification. Please insert the Substitute Sequence

Listing enclosed herewith immediately after the Abstract.

At the top of page 61, please delete "CLAIMS" and substitute
--What is claimed is--

IN THE CLAIMS:

Please cancel claims 1-23 without prejudice and without
disclaimer of the subject matter contained therein.

Please add the following claims:

--24. A method for assessing the agronomical fitness of a
plant or plant material comprising the steps of:

- a) subjecting an explant of said plant or plant material
to a stress condition;
- b) measuring the electron flow in the mitochondrial
electron transport chain to assess agronomical fitness
in cells of said explant of said plant or said plant
material;
- c) comparing said measurement to that of explants of
control plants or control plant material, under similar
conditions as for said explants of said plant or plant
material, wherein the greater the amount of electron
flow the fitter said plant or plant material.--

--25. The method of Claim 24, wherein said electron flow in the mitochondrial electron transport chain is determined by measuring the capacity of said explant subjected to said stress condition to reduce 2,3,5-triphenyltetrazolium chloride.--

--26. The method of Claim 24, wherein the electron flow in the mitochondrial electron transport chain is determined by measuring the capacity of said explant subjected to said stress condition to reduce 3-(4,5-dimethylthiazol-2-yl)-2,3 diphenyl-2H-tetrazolium.--

--27. The method of Claim 24, wherein said stress condition is selected from salt stress, osmotic stress, stress by incubation in the presence of an inhibitor of poly-ADP-ribose polymerase, stress from extreme temperatures, stress by treatment with sublethal doses of chemicals, stress by treatment with sublethal doses of herbicides, stress by treatment with sublethal doses of heavy metals and stress by irradiation with ultraviolet light.--

--28. The method of Claim 24, wherein said stress condition is salt stress.--

--29. The method of Claim 28, wherein said salt stress is induced by incubation in K-phosphate buffer comprising between 10mM and 80 mM K-phosphate.--

--30. The method of Claim 24, wherein said stress condition is osmotic stress.--

--31. The method of Claim 30, wherein said osmotic stress is induced by incubation in a buffer comprising about 2% sucrose.--

--32. The method of Claim 24, wherein said stress condition is incubation in the presence of an inhibitor of poly-ADP-ribose polymerase.--

--33. The method of Claim 32, wherein said inhibitor of poly-ADP-ribose polymerase is selected from niacinamide, picolinamide, 5-methyl nicotinamide, methylxanthine, thymidine, benzamide, 3-methoxybenzamide, 3-aminobenzamide, 2-aminobenzamide, pyrazinamide, theobromine and theophylline.--

--34. The method of Claim 32, wherein said inhibitor is present in a concentration of from about 100 mg/L to about 1,000 mg/L.--

--35. The method of Claim 24, wherein said explant is selected from callus, hypocotyl explants, shoots, leaf disks and whole leaves.--

--36. The method of Claim 24, wherein said plant or plant material is a transenic plant or transgenic plant material.--

REMARKS

Claims 24-36 are now pending in this application.

Support for new claims 24-36 may be found on pages 15-23 of the specification. Additional support for claims 33 and 34 may be found on pages 6 and 8, respectively.

The present application is a divisional of parent application Serial No. 08/817,188, filed May 15, 1997, which is filed to pursue subject matter not covered or specifically claimed in the allowed claims of the parent application.

In fulfillment of the requirements under 37 C.F.R. §§1.821-1.825 applicants respectfully request that the disk copy of the sequence listing submitted on July 22, 1999, as file 2121-127P.sub, in parent application No. 08/817,188, be transferred to the present application.

Favorable action and early allowance of the claims are

respectfully requested.


If the Examiner has any questions concerning this application, he is requested to contact the undersigned at (703) 205-8000 in the Washington, D.C. area.

If necessary, the Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account No. 02-2448 for any additional fees required under 37 C.F.R. § 1.16 or under 37 C.F.R. § 1.17; particularly, extension of time fees.


Respectfully submitted,

BIRCH, STEWART, KOLASCH & BIRCH, LLP

By


Gerald M. Murphy, Jr.
Reg. No. 28,977

Post Office Box 747
Falls Church, VA 22040-0747
(703) 205-8000


GMM/MAA/bsh
enclosure

ABSTRACT OF THE DISCLOSURE

The invention concerns a process for producing transgenic plant cells, which comprises: contacting a culture of plant cells with an inhibitor of poly-(ADP-ribose) polymerase, prior to transformation, for a period of time sufficient to reduce the response of the cultured cells to stress and to reduce their metabolism. The untransformed cells are then contacted with foreign DNA comprising at least one gene of interest under conditions in which the foreign DNA is taken up by the untransformed cells and the gene of interest is stably integrated in the nuclear genome of the untransformed cells to produce the transgenic cells. The transgenic plant cells are recovered from the culture. The invention further concerns a process for increasing the frequency of obtaining transgenic plant cells, via Agrobacterium-mediated transformation, which comprises: contacting a culture of plant cells with an inhibitor or poly(ADP-ribose) polymerase prior to transformation for a period of approximately 1 to 2 days or culturing transgenic plant cells after transformation in a medium containing an inhibitor of poly(ADP-ribose)polymerase for a period of time of approximately 1 to 14 days.

- 1 -

GENETIC TRANSFORMATION USING A PARP INHIBITOR

This invention is related to tissue culture of eucaryotic cells and improved techniques to obtain genetically transformed eucaryotic cells and organisms, such as transgenic plant cells or plants, by lowering the stress reaction of cultured eucaryotic cells prior to contacting the cells with foreign DNA, particularly by specific inhibition of poly-(ADP-ribose) polymerase.

Background to the invention

Over the years many techniques for the genetic transformation of higher organisms (animals and plants) have been developed. In these techniques it is the ultimate goal to obtain a transgenic organism, e.g. a plant, in which all cells contain a foreign DNA comprising a gene of interest (the so-called transgene) stably integrated in their genome, particularly their nuclear genome.

Transformation is a complex process which always involves the contacting of starting cells with a DNA, usually a DNA comprising foreign gene(s) of interest. The contacting of the cells with the DNA is carried out under conditions that promote the uptake of the DNA by the cells and the integration of the DNA, including the gene(s) of interest into the genome of the cell.

Starting cells for transformation are usually cells that have been cultured in vitro for some time. After contacting the cells with the DNA, the transformed cells generally need to be cultured in vitro for a certain period in order to separate the transformed cells from the non-transformed cells and, in the case of plants, to regenerate transformed plants from the transformed cells. Indeed, complete plants can be regenerated from individual transformed cells thus ensuring that all cells of the regenerated plant will contain the transgene.

- 2 -

In many plants, genetic transformation can be achieved by using the natural capacity of certain Agrobacterium strains to introduce a part of their Ti-plasmid, i.e. the T-DNA, into plant cells and to integrate this T-DNA into the nuclear genome of the cells. It was found that the part of the Ti-plasmid that is transferred and integrated is delineated by specific DNA sequences, the so-called left and right T-DNA border sequences and that the natural T-DNA sequences between these border sequences can be replaced by foreign DNA (European Patent Publication "EP" 116718; Deblaere et al, 1987 Meth.Enzymol. 153:277-293).

Certain plant species have proven to be recalcitrant to Agrobacterium mediated transformation and in these species, as well as in animals, genetic transformation has been achieved by means of direct gene transfer by which DNA is inserted into the cells by physical and/or chemical means, such as by electroporation, by treatment of the cells with polyethyleneglycol (PEG), by bombardment of the cells with DNA-coated microprojectiles, etc. (WO 92/09696; Potrykus et al, 1991, Annu.Rev.Plant Physiol.Plant Mol.Biol. 42:205-225).

Genetic transformation of eucaryotic cells is generally a random event, i.e. the transgene is integrated in the genome at random positions. Often several copies (or parts of copies) of the transforming DNA are integrated in a single position, and/or at different positions, resulting in a transformed cell containing multiple copies of the transgene.

The expression of the transgene is known to be influenced by its position in the genome. For instance, a foreign DNA, when introduced in a plant cell appears to integrate randomly in the plant genome. Examination of independently transformed plants has shown a high degree of variability (up to 100-fold) in the expression level of the introduced gene. Several studies have shown no correlation between this "between-transformant variability" and the copy

- 3 -

number of the introduced DNA at a given locus. It has been suggested that some of the variability in expression of introduced genes in transgenic plants is a consequence of "position effects" caused by influences of adjacent plant genomic DNA. Other factors that could contribute to the variability in expression are physiological variability of the plant material, differences in the number of independent T-DNA loci in different transformants or the inhibitory effects of certain T-DNA structures on gene expression. Between-transformant variability in expression has been observed for the majority of introduced genes in transgenic plants. The variability in expression of many introduced genes in independent transgenic plants necessitates large numbers of transgenic plants to be assayed to accurately quantitate the expression of the gene. It would be of great importance if the amount of between-transformant variability could be reduced (Dean et al, 1988, NAR 16:9267-9283).

If the transgene is under the control of a tissue-specific promoter, with the expectation that it will be expressed in selected tissues of the transformed organisms, the position effects can lead, at least in some transformants, to loss of specificity of the promoter and expression of the transgene in undesired tissues, e.g. in tissue cultured in vitro.

Factors that are known to influence the efficiency and quality of the genetic transformation process are the method of DNA delivery, specific tissue culture conditions, the physiological and metabolic state of the target cells etc. Direct gene transfer methods for instance are generally known to result in transformed organisms with a high copy number of the transgene.

Many of these factors are not under the control of man.

- 4 -

Summary of the Invention

This invention provides a process for producing transgenic eucaryotic cells, particularly plant cells. The process comprises contacting a culture of untransformed cells with an

5 inhibitor of poly-(ADP-ribose) for a period of time sufficient to reduce the response of the cultured cells to stress and to reduce the metabolism of the cultured cells, particularly to reduce the electron flow in the mitochondrial electron transport chain. The untransformed cells are then contacted with foreign DNA comprising at least one gene of interest under conditions in which
10 the foreign DNA is taken up by the untransformed cells and the gene of interest is stably integrated in the nuclear genome of the untransformed cells to produce the transgenic cells which are recovered from the culture.

The process may preferably comprise contacting untransformed eucaryotic
15 (e.g.) cells with foreign DNA comprising at least one gene of interest under conditions in which the foreign DNA is taken up by the untransformed cells and the gene of interest is stably integrated in the nuclear genome of the untransformed cells to produce the transgenic cells. The untransformed cells are cultured in vitro in a culture medium containing an inhibitor of poly-(ADP-
20 ribose) polymerase, preferably niacinamide, preferably for at least 2 to 3 days, particularly for at least 4 days (e.g. 4-5 days), before the contacting of the untransformed cells with the foreign DNA. The inhibitor can in addition also be applied to cultured cells that are being contacted or that have been contacted
the foreign DNA.

Description of the Invention

25 The present invention is based on the observations that poly-(ADP-ribose) polymerase (PARP) is an enzyme that is involved in regulating the general metabolic state of an eucaryotic cell and that inhibition of this enzyme can be

- 5 -

used to influence the metabolic state of cells which are targeted for transformation (or which are being transformed) to increase the efficiency and/or quality of transformation.

5 In mammals, PARP is a monomeric nuclear Zn-finger protein of about 116 kD that is closely associated with nuclear DNA, particularly with actively transcribed euchromatic regions (Shah et al, 1995, Anal.Biochem. 227:1-13). The protein is normally an inactive enzyme but is known to be activated by nicked or otherwise damaged DNA. Active PARP transfers the ADP-ribose moiety of NAD⁺ to various nuclear proteins to synthesize a polymer of ADP-ribose bound to these proteins which include PARP itself, polymerases, histones, endonuclease etc. The proteins on which such a ADP-ribose polymer is synthesized become biologically inactive (de Murcia et al, 1994, TIBS 19:172-176; Cleaver et al, 1991, Mutation Res. 257:1-18).

10 The biological function of PARP is largely unknown but the enzyme has been implicated in :

- enhancement of DNA repair (Satoh et al, 1992, Nature 356:356-358; Satoh et al, 1993, J.Biol.Chem. 268:5480-5487),
- recombination events : in general inhibition of PARP is observed to inhibit illegitimate recombination and to increase intrachromosomal recombination but it does apparently not affect extrachromosomal recombination (Farzaneh et al, 1988, NAR 16:11319-11326; Waldman and Waldman, 1990, NAR 18:5981-5988; Waldman and Waldman, 1991, NAR 19:5943-5947),
- 25 - regulation of gene expression : inhibition of PARP is observed to decrease gene expression (Girod et al, 1991, Plant Cell, Tissue and Organ Culture 25:1-12);
- reducing the amount of available NAD⁺ (and by consequence its precursor ATP) : this results in a general slowing down of cell metabolism (Lazebnik

- 6 -

et al, 1994, 371:346-347; Gaal et al, 1987, TIBS 12:129-130; Cleaver et al, supra)

It is known that PARP can be efficiently inhibited by a number of compounds (Durkacz et al, 1980, Nature 283:593-596; Sims et al, 1982, Biochemistry 21:1813-1821). Examples of such compounds are certain pyridine analogs such as nicotinamide analogues, including niacinamide, picolinamide, and 5-methyl nicotinamide; purine analogs like methylxanthines; thymidine; pyrazinamide analogs and many aromatic amides such as many benzamide analogs including benzamide, 3-methoxybenzamide and 3-aminobenzamide. For the purpose of this invention a PARP inhibitor is generally understood as any specific inhibitor of poly-(ADP-ribose) polymerase which can be taken up by a eucaryotic cell, particularly a plant cell, and which has a inhibition constant (K_i) which is lower than 1×10^{-5} , particularly lower than 1×10^{-6} . Generally it is desired that the PARP inhibitor used with this invention be a compound which in human lymphocytes, cultured in medium containing the inhibitor at a concentration of 2 mM, results in a 80-90 % inhibition of PARP (Sims et al, supra). Generally it is also preferred that cells cultured in medium containing the PARP inhibitor retain their capacity of DNA repair.

Particularly preferred PARP inhibitors are those listed above and especially niacinamide (nicotinamide), picolinamide, 5-methylnicotinamide, 2-aminobenzamide, pyrazinamide, theobromine and theophylline. Particularly niacinamide is believed to be a useful inhibitor for the purpose of this invention.

Basically the present invention provides a modification of existing procedures for the genetic transformation of eucaryotic cells, particularly plant cells, by including in the medium in which such cells are cultured a PARP inhibitor such as niacinamide, for a defined period of time. In particular the PARP inhibitor is added to the culture medium at least 1 day prior to the moment (the "contacting

- 7 -

time") at which the cells are contacted with foreign DNA comprising one or more genes of interest. However, depending on the purpose, the PARP inhibitor may also be added to the culture medium during and/or after the contacting time or even solely after the contacting time.

5 In one aspect of this invention treatment of cultured cells, tissues or organs with PARP inhibitors may be used to increase the quality of transformation as measured by the copy number of the transgene and by variation in transgene expression (quality and quantity) in the transformed cells and in organisms
10 obtained from the transformed cells.

In many conventional procedures for genetic transformation of eucaryotic cells, particularly plant cells, cultured cells, tissues or organs will be used as starting material and cells in such cultures will be contacted with foreign DNA comprising at least one gene of interest (i.e. the transgene) under conditions
15 that will promote the uptake of the foreign DNA in the cells and the ultimate integration of the foreign DNA into the genome of the cells.

In one embodiment of the invention it is preferred that a PARP inhibitor is added to the culture medium for a period of at least 2-3 days, preferably at least about 3 days, prior to contacting the cells with the foreign DNA. The exact
20 period in which the cultured cells are incubated in PARP inhibitor containing medium is believed not to be critical but should probably not exceed 4 weeks. It appears that 2-14 days, particularly 3-10 days, is an optimal period and best results were obtained with an incubation period of approximately 4 to 5 days prior to the contacting time. Generally it is believed that 4 days is a useful
25 period for the PARP inhibitor to be added to the culture medium prior to the contacting time.

The concentration of the PARP inhibitor in the medium is also believed to have an effect on the inhibition of PARP, which varies depending on the nature of the cells (species, tissue explant, general culture conditions, etc.) However,

- 8 -

within certain concentration ranges, the effect is minimal, especially when the cultured cells are not incubated for longer than 14 days. The optimal concentration range of PARP inhibitor in the medium may vary depending on the species from which the tissue, cell or cell culture is derived, but 250 mg/l (about 2 mM) is believed to be a suitable concentration for many purposes (e.g. for use with material derived from wheat). However, when nicotinamide is used in combination with plant material derived from rice, the concentration of nicotinamide should preferably be between 500 mg/l (about 4 mM) and 1000 mg/l (approx. 8 mM). On the other hand, when nicotinamide is used in combination with plant material derived from corn, the concentration of nicotinamide should preferably be 100 mg/l. Likewise, a concentration of 100 mg/l is already effective for wheat-derived plant material, but higher concentrations may be used. The optimal concentration will depend on the nature of the specific PARP inhibitor used, particularly on its strength of inhibition (as measured by its K_i and/or by its percentage inhibition of PARP under standard conditions - Sims et al, supra). It was found for instance that the optimal concentration for nicotinamide is approximately 250 mg/l (i.e. about 2 mM) but it is believed that concentrations up to 1000 mg/l (approx. 8 mM) and as low as 150 mg/l (approx. 1.25 mM), even as low as 100mg/l can be used to good effect. Preferably the nicotinamide concentration should be between 200 and 300 mg/l, i.e. between approximately 1.5 mM and 2.5 mM. In similar conditions, the optimal concentration for more potent PARP inhibitors such as 3-methoxybenzamide is about 0.5 mM, but it is believed that concentrations up to 2 mM and as low as 0.1 mM can be used to good effect. Similar concentrations apply to other PARP inhibitors.

If incubation times of longer than 14 days are used it is believed that the PARP inhibitor concentration should be reduced below 2 mM (e.g. between 0.5 mM and 1.5 mM and particularly approximately 0.8 mM).

- 9 -

For other PARP inhibitors optimal concentrations can be easily established by experimentation in accordance with this invention.

During transformation it is not known whether the integration of the DNA into the genome of the cell occurs immediately after uptake of DNA by the cell. It may very well be that the foreign DNA exists as free DNA within the cell for a certain period after the contacting time. Therefore cultured cells may be further incubated in medium containing a PARP inhibitor during and, for a limited period of time after, contacting the cells with the foreign DNA. Again the length of the incubation period is not critical but is preferably 2-10 days, particularly approximately 4 days. It is preferred that the inhibitor concentration of the PARP inhibitor in the culture medium after the contacting time should be below 2 mM, between 0.8 and 1 mM. If the cells that are to be transformed are not obtained from a cell or tissue culture (e.g. when intact tissue of an organism is contacted directly with DNA, as for example described in WO 92/09696) the PARP inhibitor may still be applied to the target cells prior to the contacting time but the addition of the PARP inhibitor to the culture of the transformed cells during or after the contacting time is preferred.

As indicated above, PARP inhibitor treatment of cultured cells for at least 2-3 days increases the quality of transformation. Indeed the number of copies of the foreign DNA is expected to be generally lower and variation in expression profile (level - i.e. the quantity - of expression as well as spatial and time distribution - i.e. the quality - of expression in the transgenic organism) of the gene(s) of interest in the foreign DNA, due to position effects, is decreased. However, at least in this aspect of the invention, the efficiency of transformation can be decreased. The efficiency of transformation as used herein can be measured by the number of transformed cells (or transgenic organisms grown from individual transformed cells) that are recovered under standard experimental conditions (i.e. standardized or normalized with respect to amount

- 10 -

of cells contacted with foreign DNA, amount of delivered DNA, type and conditions of DNA delivery, general culture conditions etc.).

Therefore it is preferred that the invention is used with transformation procedures that already have a high efficiency, such as Agrobacterium mediated transformation of dicots and direct gene transfer in monocots, particularly cereals (e.g. electroporation or particle bombardment of compact embryogenic callus in cereals - see WO 92/09696). Indeed these transformation procedures are generally highly efficient but the quality of transformation is generally poor. Position effects are large and, especially with direct gene transfer, the copy number of the transgene is often exceptionally high making analysis and selection of optimal transformants, as well as further breeding with the transformants, difficult.

In another aspect of this invention treatment of cultured plant cells for a short period of time (i.e. 1 day to maximally 2 days) prior to, or after contacting the cells with DNA may be used to increase the efficiency of Agrobacterium mediated transformation of plant species, such as many monocots, particularly the major cereals such as wheat and corn, for which this method is generally inefficient. It is believed that treatment of cultured plant cells during the contacting time may result in a lower transformation efficiency, and might therefore not be suitable for this aspect of this invention. Likewise, it is believed that for the purpose of this aspect of the invention, the optimal treatment with a PARP inhibitor is 1 day to maximally 2 days prior to the contacting time, or alternatively 1 to maximally 2 days after the contacting time. In this embodiment of the invention the contacting of the plant cells with the DNA should of course be understood as contacting the cells with an appropriate Agrobacterium strain harboring an artificial T-DNA containing the foreign DNA with the gene(s) of interest. In this embodiment of the invention the quality of transformation is expected not to be affected but this is generally deemed to be

- 11 -

of lesser importance since Agrobacterium mediated transformation, being a biological process, already results in a generally low copy number of the transgene in the transformed plant cells.

5 In accordance with this invention the addition of PARP inhibitors, such as niacinamide, to the culture medium of eucaryotic cells, can be used in combination with any known transformation procedure that requires cells, tissues or organs cultured in vitro as starting cells to be contacted with foreign DNA. The process of this invention is thus generally identical to existing
10 conventional transformation methods except for the fact that at some times during the tissue culture of the cells, a PARP inhibitor is added to the culture medium.

The cell of a plant, particularly a plant capable of being infected with Agrobacterium such as most dicotyledonous plants (e.g. Brassica napus) and
15 some monocotyledonous plants, can be transformed using a vector that is a disarmed Ti-plasmid containing the gene(s) of interest and carried by Agrobacterium. This transformation can be carried out using conventional procedures (EP 0,116,718; Deblaere et al, supra; Chang et al, 1994, The Plant Journal 5:551-558). Preferred Ti-plasmid vectors contain the foreign DNA
20 between the border sequences, or at least located to the left of the right border sequence, of the T-DNA of the Ti-plasmid. Of course, other types of vectors can be used to transform the plant cell, using procedures such as direct gene transfer (as described, for example, in EP 0,233,247), pollen mediated transformation (as described, for example, in EP 0,270,356, PCT patent publication "WO" 85/01856, and US patent 4,684,611), plant RNA virus-
25 mediated transformation (as described, for example, in EP 0,067,553 and US patent 4,407,956) and liposome-mediated transformation (as described, for example, in US patent 4,536,475). Cells of monocotyledonous plants such as the major cereals including corn, rice, wheat, barley, and rye, can be

- 12 -

transformed (e.g. by electroporation) using wounded or enzyme-degraded intact tissues capable of forming compact embryogenic callus (such as immature embryos in corn), or the embryogenic callus (such as type I callus in corn) obtained thereof, as described in WO 92/09696. In case the plant to be transformed is corn, other recently developed methods can also be used such as, for example, the method described for certain lines of corn by Fromm et al., 1990, Bio/Technology 8:833; Gordon-Kamm et al., 1990, Bio/Technology 2:603 and Gould et al., 1991, Plant Physiol. 95:426. In case the plant to be transformed is rice, recently developed methods can also be used such as, for example, the method described for certain lines of rice by Shimamoto et al., 1989, Nature 338:274; Datta et al., 1990, Bio/Technology 8:736; and Hayashimoto et al., 1990, Plant Physiol. 93:857; Hiei et al, 1994, The Plant Journal 6:271-282).

The transformed cell can be regenerated into a mature plant and the resulting transformed plant can be used in a conventional breeding scheme to produce more transformed plants with the same characteristics or to introduce the gene(s) of interest in other varieties of the same related plant species. Seeds obtained from the transformed plants contain the chimeric gene(s) of this invention as a stable genomic insert. Thus the gene(s) of interest when introduced into a particular line of a plant species can always be introduced into any other line by backcrossing.

In animals pluripotent embryonic or somatic stem cells can be used as target for transformation (Capecchi et al, 1989, TIG:5:70-76).

The transformed cells and organisms of any plant or animal species, produced by the process of this invention, contain the foreign DNA as a stable insert in their genome, particularly in regions of the genome that remain transcriptionally active in the untransformed cells that have been exposed to a PARP inhibitor in

- 13 -

accordance with this invention. As described above it is believed that in cells treated with a PARP inhibitor for at least 3 days, particularly for at least 4 days, only a limited number of genomic regions will remain transcriptionally active. In this regard the transformed cells, obtained with this process of the invention, will be characterized by having the foreign DNA integrated in a limited number of genomic regions. That the transformed cell or organism was obtained by this process of the invention can thus be easily ascertained by 1) culturing transformed cells or tissues under conditions that are similar as those in which the untransformed cells or tissues were grown or incubated prior to the integration of the foreign DNA in the genome (i.e. incubating in medium containing 250 mg/l niacinamide for 4-5 days prior to the contacting time), and 2) monitoring the expression of at least one transgene in the foreign DNA that is expected to be expressed under normal tissue culture conditions (i.e. a selectable marker gene under the control of a promoter that directs expression in tissue culture). Under the above conditions the transformed cells or tissues of this invention express the relevant transgene in the tissue culture at essentially the same levels whether or not a PARP inhibitor is present in the culture medium. It is thus expected that, for instance after 4-5 days of culturing of the transformed cells in medium containing the PARP inhibitor, mRNA levels are not significantly decreased, i.e. do not become lower than 75%, preferably not become lower than 90%, when compared to the mRNA levels observed in cells cultured in medium not containing the inhibitor. Indeed, if the relevant transgene is integrated in other regions of the genome (i.e. in regions that are normally not transcriptionally active in cells treated with PARP inhibitor according to this embodiment of the invention), the expression of the relevant transgene is considerably reduced after incubation of the cells in medium containing the PARP inhibitor for at least 3 days, e.g. 4-5 days (i.e. mRNA levels will drop below 75%, particularly below 50%, more particularly below

- 14 -

30%) when compared to the mRNA levels observed in cells cultured in medium not containing the inhibitor).

The method of the present invention can in principle be used to transform eucaryotic cells with any foreign DNA. Generally the foreign DNA comprises at least one gene of interest comprising 1) a promoter region with a promoter capable of directing transcription of DNA into a RNA in cells of the eucaryotic, e.g. plant, species that is to be transformed and 2) a coding region coding for a RNA or protein. Most often the gene of interest will also comprise 3) a 3' untranslated region of a eucaryotic gene containing a polyadenylation signal.

The promoter can be selected to direct expression in selected tissues of the eucaryotic organism. Such a tissue-selective promoter is not expected to direct expression in other non-selected tissues. For instance promoters are known that direct expression selectively in stamen tissues of a plant and such promoters have been used to produce male sterile plants and other plants useful for producing hybrids (EP 344029; EP 412911; WO 9213956; WO 9213957; Mariani et al, 1990, Nature 347:737-741; Mariani et al, 1992, Nature 357:384-387).

It is believed that the method of the present invention is particularly useful to transform eucaryotic cells with at least one gene of interest comprising a tissue-selective promoter, such as a stamen selective promoter, especially if expression of the gene of interest in the organism, such as a plant, outside the selected tissues (where the tissue-selective promoter is active, i.e. directs expression) is undesired for example because the gene product (for instance a protein such as a ribonuclease, e.g. barnase) is capable of killing or disabling the cells in which they are produced. In such cases expression of the gene of interest in tissue culture, or in non-selected tissues of the organisms can negatively affect the quality as well as the apparent efficiency of transformation. When the method of this invention is used, the overall efficiency of

- 15 -

transformation may be reduced but the average quality of transformation is expected to be significantly improved because of lower copy number of the gene of interest in the genome of the transformed cells and because of reduced position effects i.e. the general integration of the gene of interest in the genomes at locations that minimally affect the transcriptional properties of the promoter of the transgene.

The foreign DNA used in the method of this invention generally also comprises a selectable marker gene the expression of which allows the selection of transformed cells (or organisms) from non-transformed cells (or organisms).

Such selectable marker gene generally encodes a protein that confers to the cell resistance to an antibiotic or other chemical compound that is normally toxic for the cells. In plants the selectable marker gene may thus also encode a protein that confers resistance to a herbicide, such as a herbicide comprising a glutamine synthetase inhibitor (e.g. phosphinothricin) as an active ingredient. An example of such genes are genes encoding phosphinothricin acetyl transferase such as the sfr or sfrv genes (EP 242236; EP 242246; De Block et al, 1987 EMBO J 6:2513-2518).

The inventors also found that the initial reaction of cells, particularly cells contacted with PARP inhibitors, is a stress reaction which enhances free radical production by the cell. However, this stress only lasts for a limited period of time after which further contact with the PARP inhibitor causes a decrease in cell metabolism, particularly a decrease in electron flow in the mitochondrial electron transport chain. Therefore, the invention also relates to a new method to assess the agronomical fitness of a population of transformed plants to determine in which lines the plants have a foreign DNA integrated in their genomes in a way that agronomical performance is not or substantially not affected. The assay is based on comparative reaction of transgenic cells and corresponding untransformed controls to stress conditions.

- 16 -

The method comprises exposing the transgenic cells to stress conditions which induce the production of free radicals in the tissues or the cells, measuring the amount of free radicals produced in the transgenic cells with the amount of free radicals produced in control cells exposed to similar stress conditions. Preferably the cells of the transgenic organism to be assayed are exposed to stress conditions by being treated with a substance which induces increasing osmotic and/ or salt stress on the cells.

The properties of PARP inhibitors, such as niacinamide, to enhance free radical production in cells incubated with the inhibitor for not longer than 2 days, preferably not longer than 1 day, can be used to assay the (relative) fitness of a population of transgenic eucaryotic organisms, particularly plants.

The term fitness used herein is intended to designate the agronomical performance of a population of plants, as measured for instance by its yield (e.g. its seed yield) as compared to a given reference population. Agronomical performance is generally thought to be correlated with the general resistance of the plants to a range of stress conditions which are likely to be encountered in the field locations where the plants are normally grown. For any population of transformed plants (i.e. a transgenic line) the relevant reference population is a population of untransformed plants of the same variety.

It is known that in transformed plants and other organisms transgene expression may be qualitatively and quantitatively influenced by the genomic domain in which the transgene(s) are integrated, that undesired transgene expression may interfere with cell metabolism (e.g. when the transgene encodes a cytotoxic protein), that mutations may be induced in the transformed organism either by somaclonal variation or by insertional inactivation of

- 17 -

endogenous genes by the transgene(s), or that expression of endogenous genes may be deregulated by sequences in the foreign DNA. As a consequence many transformed lines may not be agronomically useful.

The assay of this invention will for example allow to identify a line (i.e. a group of genetically similar plants) of transformed plants that have the transgene(s) integrated in regions that minimally affect the fitness of the plants, thus avoiding the extensive laboratory, greenhouse and/or field evaluations which are normally required to identify the transformants with the best agronomical properties.

The assay in accordance with this invention essentially comprises the incubation of cells or tissues of transformed plants of a particular transgenic line (e.g. callus, hypocotyl explants, shoots, leaf disks, whole leaves etc.) preferably with a PARP inhibitor (although for some plant species this is not necessary) under a range of conditions which induce the production of a different amount of free radicals in the tissues. An incubation time of approximately one day is normally sufficient to generate the desired amount of free radicals. Appropriate controls, i.e. untransformed tissues obtained from untransformed plants at the same developmental stage and grown in the same conditions as the transformed plant from which the transformed tissue was obtained, are subjected to the same treatment. Preferably the untransformed line is identical to the transgenic line except for the presence of the transgene(s).

For each plant line (control or transformant) it is preferred that a number of plants is assayed.

Useful conditions for the incubation of the untransformed and transformed tissues are those which induce increasing osmotic and salt stress in the incubated cells or tissues. For example a series of buffers with different salt

- 18 -

concentrations containing a PARP inhibitor can be made. A useful buffer series is a K-phosphate buffer containing 2% sucrose and 250 mg/l niacinamide in which the K-phosphate concentration is increased from anywhere between 10 to 80 mM (e.g. in steps of 5 mM, i.e. 10, 20, 25, 30, 35, 40, 45, 50, 55, 60 mM). The K-phosphate concentrations will induce mild but increasing salt and osmotic stress in plant cells. The niacinamide in the medium further enhances radical production and stress on the plant cells. The range of K-phosphate concentrations used will depend on the natural sensitivity of the plant species (or if desired the plant line) to the salt and osmotic stress. In sensitive plant species, which will not tolerate high salt stress, the maximum K-phosphate concentration may for instance be 50 mM, in less sensitive species this maximum K-phosphate concentration can be increased up to 70 or 80 mM or even higher. For each plant species the minimum and particularly the maximum salt (e.g. K-phosphate) concentration can be determined experimentally for an untransformed line - the only requirement is that at all concentrations used the plant tissue remains viable. Although the addition of a PARP inhibitor to the medium, such as niacinamide, is preferred it is not required for assaying plant species that are very sensitive to salt and/or osmotic stress.

After the one day incubation the capacity of the transformed and control tissues to reduce 2,3,5-triphenyltetrazolium chloride (TTC) is measured e.g. by the following procedure which is modified from Towill and Mazur (supra):

- incubate the tissues for 1 to 4 hours in K-phosphate buffer (pH 7.4) containing 10 mM TTC and 0.1 % Tween20. As a control similar plant material is incubated in the same buffer without TTC.
- extraction of reduced TTC (e.g. freezing at -70°C followed by thawing at 40°C and shaking the plant material in ethanol for 45-60 minutes)

- 19 -

- spectrophotometric quantification of reduced TTC at 485 nm (optical density OD₄₈₅; for chlorophyll poor plant material) or 545 nm (OD₅₄₅; for chlorophyll rich plant material). The O.D. of the control extract is subtracted from the OD of the TTC-reacted extracts. In the above conditions 0.1 mM reduced TTC corresponds to an OD₄₈₅ of 0.214 or OD₅₄₅ of 1.025 (light path 1 cm).
- the reducing capacity of the transformed plant line is compared to that of the control line.

The amount of reduced TTC is determined by the intensity of the cytochromal and alternative respiratory pathways and the radical concentration in the tissues which, in turn are determined by the presence of mutations, the expression of genes affecting the metabolic activity of the plant cells, the developmental stage and the reaction of the tissue to external factors, such as stress factors.

The TTC reducing capacity (as for instance measured by the O.D. at 485 nm) for tissues incubated at high salt concentration (TTC-high) is expressed as the percentage of the TTC reducing capacity of the tissues incubated at a low salt concentration (TTC-low); in other words a TTC-ratio value is calculated as follows:

$$\text{TTC-ratio} = \text{TTC/high} \cdot 100 / \text{TTC/low}.$$

The value of TTC-ratio is a measure of the fitness of a plant line as compared to a control line.

The determination of TTC-low and TTC-high will depend on the sensitivity of the plant species to the applied salt stress. Usually TTC-low will correspond to a salt concentration between 10 and 25 mM K-phosphate, e.g. at 20 mM while TTC-high will correspond to a salt concentration between 50 and 80 mM K-phosphate. The only requirement is that TTC-high should be significantly lower than TTC-low; preferably TTC-high should be lower than 50% of TTC-low, particularly lower than 30% of TTC-low. For instance for Brassica napus, TTC-

- 20 -

low and TTC-high can be typically obtained from tissues incubated at respectively 20mM and 60 mM K-phosphate buffer containing 250 mg/l niacinamide. TTC-high and TTC-low, for both the transformed and untransformed line, will usually be an average obtained from several measurements taken on a number of tissue explants from a number of plants of each line. For instance for each line of Brassica napus about 32 leaf discs (diameter 1 cm) from 8 different plants (i.e. about four leaf discs per plant) can be assayed to determine 32 TTC-high and 32 TTC-low values which are averaged to obtain the TTC-high and TTC-low values used for the calculation of TTC-ratio. Other examples of sample sizes which have been used are 35 shoots from Arabidopsis thaliana, or 150 hypocotyl explants derived from about 25 seedlings of Brassica napus.

Transformed lines with a value of TTC-ratio which does not deviate more than 20%, preferably not more than 10% of the TTC-ratio value of the control line are selected. These lines are likely to have the transgene(s) integrated in regions that minimally affect the fitness of the plants.

It is clear that additional information considering the fitness of the plant material studied can be obtained by comparing the TTC-reducing capacity of the plant material in absence of a PARP-inhibitor with the TTC-reducing capacity of the plant material in the presence of a PARP-inhibitor for each experimental point of the buffer series mentioned above.

While the TTC-reduction assay is especially suitable for the identification of transgenic plants, where transgenes are integrated in regions that minimally affect the fitness of the plants, this test can also be successfully applied to discriminate mutant plants, cells or cell lines from the wild-types.

- 21 -

The TTC-reducing assay can further be used in a modified way to determine the quality and the fitness of plant material, for example plant material to be used in transformation experiments (i.e. whether particular plant material, e.g. explants, is suitable as starting material). To this end the TTC-reducing assay can be adapted for example in the following way:

1. A sample of the plant material to be tested for its suitability for transformation, is incubated for one day in plant culture medium or a buffer containing 2% sucrose and a K-phosphate concentration ranging between 10 and 80 mM, typically around 25 mM, to which a suitable amount of a PARP inhibitor, such as niacinamide has been added. For niacinamide, a preferred concentration to be used is 250 mg/L, although concentrations as low as 100 mg/L and as high as 1000 mg/L may be used. A comparable control sample of the same plant material is incubated under similar conditions without PARP inhibitor.
2. After one day of incubation the capacity of the plant material incubated with PARP inhibitor and the control plant material to reduce TTC is measured by the procedure described above.

The TTC reducing capacity (as for instance measured by the O.D. at 485 nm) for plant material incubated with PARP inhibitor (TTC-INH) is compared with the TTC reducing capacity of the control plant material incubated without PARP inhibitor (TTC-CON) and a ratio (E) is calculated as follows:

$$E = \text{TTC-INH} / \text{TTC-CON}$$

The value E is a measure of the quality and fitness of the plant material, for example explants to be transformed. It is believed that those tissues, wherein the E value is larger than or equals 1, are healthy tissues, which are particularly suitable as starting material for transformation.

The modified TTC-procedure thus allows to select those types of (cultured) plant material especially appropriate for use in a transformation procedure,

- 22 -

particularly the procedures of this invention that include the use of a PARP inhibitor.

5 As the quality of plant material will also be affected by the particular culture conditions used prior to transformation (especially cells, tissues or explants derived from plants recalcitrant to transformation) the assay of this invention is further useful to identify suitable culture conditions to obtain suitable starting plant material. Thus it has been found by the inventor that, when culturing plant material from corn, it is preferred to include proline, preferably at a
10 concentration of about 8mM, simultaneously with the PARP inhibitor, in the culture medium.

As already mentioned, incubation of cells or tissues in the presence of a PARP inhibitor for longer than 1 to 2 days leads to a general reduction in cell metabolism, particularly a reduction in the electron flow in the mitochondrial electron transport chain (after the initial increase, characteristic of healthy cells or tissues, during the first day). The period of time required to reduce the metabolism to an optimal level (for the purpose of improving the qualitative aspect of transformation) is that period after which a decrease in TTC-reducing capacity between 20% and 50%, preferably between 30 % and 40%, particularly about 35%, is achieved for plant material incubated with a PARP inhibitor (e.g. niacinamide) when compared to control plant material incubated without the PARP inhibitor (i.e. the period after which the E value is between 0.5 and 0.8, preferably is between 0.6 and 0.7, particularly is about 0.65).
20

25 It is clear that the assays of this invention can be readily adapted by one skilled in the art of the field, for example to suit the needs of the particular cell type, tissue or explant or of the particular species from which the cells, tissues or explants are derived. Furthermore the assay can be adapted to assay a

- 23 -

peculiar aspect of fitness of cells, tissue, explant or organism. For instance, it is possible to apply a type of stress different from osmotic or salt stress, such as stress brought about by extreme temperatures, by sublethal treatment with chemicals (e.g. herbicides, heavy metals) or by irradiation with UV. Furthermore, other types of PARP inhibitors, as mentioned before may be used, within the indicated concentration ranges. Although it is believed that for the purpose of the assays defined here, TTC is the most suited substrate, other indicator molecules, such as MTT (3-(4,5-dimethylthiazol-2-yl)-2,5 diphenyl-2H-tetrazolium) can be used to measure the electron flow in the mitochondrial electron transport chain downstream of the "ubiquinone pool".

Unless otherwise indicated all experimental procedures for manipulating recombinant DNA were carried out by the standardized procedures described in Sambrook et al., 1989, "Molecular Cloning: a Laboratory Manual", Cold Spring Harbor Laboratory, and Ausubel et al, 1994, "Current Protocols in Molecular Biology", John Wiley & Sons.

The polymerase chain reactions ("PCR") were used to clone and/or amplify DNA fragments. PCR with overlap extension was used in order to construct chimeric genes (Horton et al, 1989, Gene 77:61-68; Ho et al, 1989, Gene 77:51-59).

All PCR reactions were performed under conventional conditions using the VentTM polymerase (Cat. No. 254L - Biolabs New England, Beverley, MA 01915, U.S.A.) isolated from Thermococcus litoralis (Neuner et al., 1990, Arch.Microbiol. 153:205-207). Oligonucleotides were designed according to known rules as outlined for example by Kramer and Fritz (1968, Methods in Enzymology 154:350), and synthesized by the phosphoramidite method (Beaucage and Caruthers, 1981, Tetrahedron Letters 22:1859) on an applied Biosystems 380A DNA synthesizer (Applied Biosystems B.V., Maarssen,

- 24 -

Netherlands). In the examples MS medium means Murashige and Skoog medium (Murashige and Skoog, 1962, *Physiol. Plant* 15:473-479).

In the following examples, reference will be made to the following sequence listing and figures:

Sequence Listing

SEQ ID NO 1 : T-DNA of plasmid pTHW107

SEQ ID NO 2: plasmid pTS172

SEQ ID NO 3: PT72 promoter contained in plasmid pTS772

SEQ ID No 4 : plasmid pVE136

SEQ ID No 5 : T-DNA of plasmid pTHW142

Examples

Example 1 : Tissue culture of wheat embryogenic callus and Brassica napus hypocotyl explants in media containing a PARP inhibitor.

Wheat embryogenic callus was cultured on W2 medium (see Example 2). When niacinamide was added as PARP-inhibitor to the medium at a concentration of 250 mg/l (approx. 2 mM) it was observed that after 4 days the growth of the tissue was slowed down considerably (to approximately 30% of the normal rate after 4 weeks) but the tissue remained viable for extended periods of time (i.e. at least one month). If niacinamide was subsequently removed from the medium the tissue started to grow normally again. It was also observed that after 4-5 days of incubation of the plant tissue with niacinamide, the TTC-reducing capacity (Towill and Mazur, 1975, *Can J.Bot.* 53:1097-1102) of the tissue was substantially decreased probably indicating a reduction of the production of free radicals and decreased mitochondrial electron transport.

- 25 -

Similar observations were made when Brassica napus hypocotyl explants were cultured on A5 medium (see Example 3) containing 250 mg/l niacinamide. It was also observed that, in Brassica napus tissue cultured on medium containing niacinamide, no anthocyanin was produced; normally anthocyanin in tissue culture is produced in stress conditions. In addition it was observed that after 4-5 days of incubation of the plant tissue with niacinamide, the concentrations of hydroxyl free radical and dehydroascorbate in the explants were drastically decreased.

It was also observed that, after a 4 day incubation in niacinamide containing medium, the percentage of cultured cells that were in G2 phase of the cell cycle was considerably increased (up to 45 % of all cells in the culture).

The above observations are interpreted as indicating that treating cultured cells with a PARP inhibitor for about 4-5 days generally results in :

- 1) a significant reduction of the response of the cultured cells to stress as measured for instance by free radical and/or anthocyanin production , and
- 2) a reduction of the general metabolism of the cultured cells to a very basic level as indicated by the fact that the tissue growth was slowed down, and the TTC reducing capacity was decreased while the tissue remained viable.

It is inferred that under these conditions many genes in cells (e.g. cultured cells) that would normally be switched on in response to stress (such as during transformation conditions) will in fact no longer be induced. It is expected that in such cells which only display a very basic metabolism, mainly general "housekeeping genes", i.e. genes that are expressed in any cell irrespective of its differentiated state or metabolic or physiological condition, are expressed.

As it is believed that foreign DNA is preferably inserted in portions of the genome that are transcriptionally active it follows that treatment with PARP inhibitors will condition eucaryotic cells to incorporate any foreign DNA

- 26 -

preferentially in genomic regions which are transcribed in all cells and not in regions of the genome which would only be transcribed under certain conditions, i.e. stress conditions, or during differentiation. This means that the number of locations in which foreign DNA will be integrated, and the concomitant variation in expression profile of the transgene(s), will be reduced. It is further believed that this will enhance integration of foreign genes of interest in such locations which in turn will result in a more reliable and faithful expression of these genes which will be less affected by cell differentiation or cell physiological and biochemical changes due to for instance environmental conditions.

Example 2 : Transformation of wheat with a barnase gene under the control of a stamen-specific promoter using the particle bombardment

The Wheat Spring variety Pavon is grown in a greenhouse or conditioned room at 23-24°C during daytime and 18-20°C at night, with a photoperiod of 16 hours light and 8 hours dark. Developing seeds (white-greenish with white semi-liquid endosperm) were harvested, sterilized by incubation for 1 minute in 70% ethanol followed by 15 minute incubation in 1.3% NaOCl+ 0.1% Tween 20, and washed with sterile water. The sterilized seeds were either used directly or were stored for one day at 4-7°C.

Immature embryos of about 1 mm in size were isolated and were placed, with the scutellum upwards, on callus inducing medium W1 (MS medium supplemented with 3% sucrose, 40 mg/l adenine.SO₄, 0.5 mg/l thiamine.HCl, 0.5 g/l 2-[N-Morpholino] ethane sulfonic acid (Mes) pH 5.8, 0.5% agarose, 0.5 to 2.5 mg/l CuSO₄.5H₂O, 25 mg/l acetylsalicylic acid and 2 mg/l 2,4-dichlorophenoxyacetic acid (2,4-D)) and were incubated for 3 weeks at 27°C in the dark.

Embryogenic sections of the developing callus were isolated, placed on callus maintenance medium W2 (W1 medium but without acetylsalicylic acid and with

- 27 -

only 0.5 mg/l $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ and 1 mg/l 2,4-D), and incubated for 3 weeks at 24-25°C in the light (approx. 20 mEinstein/s/m² (with a photoperiod of 16 hours light and 8 hours dark).

About 2 weeks prior to bombardment the calli were cleaned up by removal of non-morphogenic (i.e. the nonembryogenic and nonmeristematic) parts and were subcultured on W2 medium.

For bombardment the calli were divided into small pieces with an average maximum diameter of about 2-3 mm. These pieces were placed at the center of a 9 cm Petridish containing W2 medium in a circle with a diameter of approx. 0.5 cm. When required niacinamide (250 mg/l) was added to the W2 medium and the tissue pieces were maintained under these conditions for 4 days after they were bombarded.

Bombardment was carried out using the Biolistic PDS-1000/He apparatus (Bio-Rad). Preparation of the microcarriers (0.4-1.2m) and the coating of the microcarriers with DNA was essentially carried out according to the manufacturer's instructions. The Petridishes containing the calli were placed at level 2 of the apparatus and the bombardment was done at 1550 psi.

For the transformation experiments the following plasmid DNA was used.

- plasmid pVE136, the sequence of which is given in SEQ ID No 4. This plasmid contains the following chimeric genes:

- P35S-bar-3'nos
- PCA55-barnase-3'nos

in which P35S is the 35S promoter of the Cauliflower Mosaic virus, bar is a DNA encoding phosphinothricin acetyltransferase (EP 242236), 3'nos is the 3' untranslated end of the Agrobacterium T-DNA nopaline synthase gene, PCA55 is a stamen-specific promoter from corn gene CA55 (WO

- 28 -

9213957), and bamase is a DNA encoding bamase (Hartley, 1988, J.Mol.Biol.202:913-915)

- plasmid pTS172 the sequence of which is given in SEQ ID No 2. This plasmid contains the following chimeric genes:

- P35S-bar-3'g7
- PE1-bamase-3'nos

in which in which P35S is the 35S promoter of the Cauliflower Mosaic virus, bar is a DNA encoding phosphinothricin acetyltransferase (EP 242236), 3'g7 is the 3' untranslated end of the Agrobacterium T-DNA gene 7, PE1 is a stamen-specific promoter from rice gene E1 (WO 9213956), bamase is a DNA encoding bamase (Hartley, 1988, J.Mol.Biol.202:913-915), and 3'nos is the 3' untranslated end of the Agrobacterium T-DNA nopaline synthase gene,

- plasmid pTS772 which is identical to pTS172 except that the region between nucleotides 2625-4313 of pTS172, containing PE1, is replaced by the sequence of SEQ ID No 3 containing the PT72 promoter. Thus, plasmid pTS772 contains the following chimeric genes:

- P35S-bar-3'g7
- PT72-bamase-3'nos

in which PT72 is a stamen-specific promoter from rice gene T72 (WO 9213956)

The bombarded calli were transferred to selective medium W2 containing 2.5 mg/l phosphinothricin (PPT) and, if necessary, 100 mg/l niacinamide. The calli that were placed on medium containing niacinamide were transferred after 4 days to niacinamide-free W2 medium containing 2.5 mg/l PPT. The cells were cultured at 24-25°C.

After two weeks the calli were subcultivated on W2 medium and after a further two weeks the growing parts of the calli were transferred to regeneration medium W4 (W1 medium but without acetylsalicylic acid and with only 0.5 mg/l

- 29 -

CuSO₄.5H₂O and 0.5 mg/l 2,4-D). Calli were subcultivated every two weeks at which time the nonmorphogenic parts of the calli were removed. When the calli started to form shoots they were transferred to W5 medium (W1 medium with half concentrated MS medium and only 0.5 mg/l CuSO₄.5H₂O and without acetylsalicylic acid and 2,4-D, but supplemented with 50 mg/l myo-inositol, 0.25 mg/l pyridoxine.HCl and 0.25 mg/l nicotinic acid) containing 2.5 mg/l PPT. For the rest of the procedure temperature was maintained at a maximum of 24°C. The calli were subcultivated every 3-4 weeks. Once the shoots started to elongate and small roots started to form, the whole calli (or if possible individual shoots) were transferred to 1 liter vessels with W6 medium (half-concentrated MS medium supplemented with 1.5% sucrose, 50 mg/l myo-inositol, 0.25 mg/l pyridoxine.HCl, 0.25 mg/l nicotinic acid, 0.5 mg/l thiamine.HCl, 0.7% agar (Difco) pH 5.8 and 0.5 mg/l CuSO₄.5H₂O) containing 2.5 mg/l PPT. Once the shoots and roots had grown out, individual shoots were separated from each other and transferred to 1 l vessels containing W6 medium with 2.5 mg/l PPT. Well developed shoots are tested for PPT resistance by means of the TLC assay (De Block et al, 1987, EMBO 6:2513-2518) or by direct assay of ammonium production in the tissue (see e.g. De Block et al, 1995, Planta 197: 619-626). Transformed shoots were finally transferred to the greenhouse into soil.

For analysis of the results the transformed plants could be subdivided according to the niacinamide treatment of the parent calli during tissue culture. Thus the following groups were distinguished:

<u>Group</u>	<u>Niacinamide treatment</u>
None	No treatment
Before 100	100 mg/l niacinamide for four days prior to bombardment
Before 250	250 mg/l niacinamide for four days prior to bombardment

- 30 -

Before/After 250 mg/l niacinamide for four days prior to bombardment
plus 100 mg/l niacinamide for four days after
bombardment

5 Results of the experiments are presented in Tables 1, 2 and 3. Plants could be obtained only from bombarded calli that were treated with niacinamide.

For the plants that were transformed with plasmid pTS172 it was demonstrated that the foreign DNA, comprising the chimeric PE1-barnase-3'nos and P35S-bar-3'g7, was stably incorporated in the wheat genome in 2 to 3 copies on the average. The fact that variation in expression profile (e.g. tissue-specificity) of the transgenes, especially the chimeric barnase genes, was decreased in transformed cells was evident from the fact that male-sterile plants that otherwise looked completely healthy could be obtained only from bombarded calli treated with niacinamide. It is believed that this is due to a more faithful expression characteristics (i.e. lack of expression) of the integrated stamen-selective barnase gene in these calli and shoots regenerated from these calli. In the control calli undesired expression of the barnase gene in tissue cultured cells might have prevented recovery of any transformed plants from these calli. It is expected that to obtain the same number of male-sterile wheat plants from control calli a much larger number of calli would have to be bombarded.

10
15
20

- 31 -

Results of wheat transformation experiments**Table 1 :**

Plasmid pTS172				
Treatment	Nr of bombarded calli	Nr of PPT- resistant calli recovered	Nr of PPT resistant plants recovered	Nr of MS plants recovered
None	60	30	1^{a)}	0
Before 250	125	30	3	3^{b)}

a) This plant proved to be fertile and to be transformed only with the chimeric bar gene

b) The obtained plants looked healthy and tillered vigorously

- 32 -

Table 2:

Plasmid pTS772				
Treatment	Nr of bombarded calli	Nr of PPT- resistant calli recovered	Nr of PPT resistant plants recovered	Nr of MS plants recovered
None	250	22	0	0
Before 250	210	75	7	3^{a)b)}
Before/ After	210	45	6	3^a

a) The obtained plants looked healthy and tillered vigorously

b) Only six plants could be analyzed for MS phenotype since one of the plants died prematurely.

- 33 -

Table 3:

Plasmid pVE136			
Treatment	Nr of bombarded calli	Nr of PPT resistant plants recovered	Nr of MS plants recovered
None	200	1	0
Before 100	800	8^{a)}	8

a) The obtained plants looked healthy and tillered vigorously

- 34 -

Example 3: Transformation of oilseed rape with a barnase gene under the control of a stamen-specific promoter using Agrobacterium mediated transformation.

Hypocotyl explants of Brassica napus were obtained , cultured and transformed essentially as described by De Block et al, 1989, Plant Physiol. 914:694-701 except for the following modifications:

- hypocotyl explants were precultured for 3 days on A2 medium (MS, 0.5 g/l Mes (pH 5.7), 1.2% glucose, 0.5% agarose, 1 mg/l 2,4-D, 0.25 mg/l naphthalene acetic acid (NAA), 1 mg/l 6-benzylaminopurine (BAP)), and then transferred to the A2 medium with or without niacinamide for another 4 days.
- infection medium A3 was MS, 0.5 g/l Mes (pH 5.7), 1.2% glucose, 0.1 mg/l NAA, 0.75 mg/l BAP, 0.01 mg/l giberellinic acid (GA3)
- selection medium A5 was 0.5 g/l Mes (pH 5.7), 1.2 % glucose, 40 mg/l adenine.SO₄, 0.5 g/l polyvinyl-polypyrrolidone (PVP), 0.5% agarose, 0.1 mg/l NAA, 0.75 mg/l BAP, 0.01 mg/l GA3, 250 mg/l carbenicillin, 250 mg/l triacillin, 5 mg/l AgNO₃.
- regeneration medium A6 was MS, 0.5 g/l Mes (pH 5.7), 2% sucrose, 40 mg/l adenine.SO₄, 0.5 g/l PVP, 0.5% agarose, 0.0025 mg/l BAP, 250 mg/l triacillin.
- healthy shoots were transferred to 1 liter vessels containing rooting medium which was either A8 or A9; A8 consists of 100-130 ml half concentrated MS, 1% sucrose (pH 5.0), 1 mg/l isobutyric acid (IBA), 100 mg/l triacillin added to 300 ml perlite (final pH 6.2); A9 consists of half concentrated MS, 1.5% sucrose (pH 5.8) solidified with agar (0.6%)

Hypocotyl explants (with or without niacinamide treatment) were infected with Agrobacterium tumefaciens strain C58C1Rif carrying T-DNA vector pTHW107 and a helper Ti-plasmid pMP90 (Koncz and Schell, 1986, Mol.Gen.Genet. 204:383-396)(or a derivative thereof).

- 35 -

Plasmid pTHW107 is a vector carrying a T-DNA comprising the following chimeric genes :

- PTA29-barnase-3'g7
- PSSU-bar-3'nos

in which PTA29 is the promoter of the TA29 gene of tobacco (EP 344029) and PSSU is the promoter of the gene of Arabidopsis thaliana encoding the small subunit of Rubisco. The complete sequence of the T-DNA of pTHW107 is presented in SEQ ID No 1.

Where required niacinamide (250 mg/l) was added to the media for the last 4 days prior to infection with Agrobacterium. Plants regenerated from transformed calli obtained on niacinamide cultured cells were observed to have a low copy number as well as to display less variation in the expression profile of the transgenes (results summarized in Table 4). Five plants regenerated from the calli obtained by transformation including niacinamide and five plants regenerated from the calli obtained by conventional transformation without niacinamide inclusion, were analyzed by Southern hybridization to determine the copy number of the transgenes, and were further analyzed for reproductive phenotype. In the non-treated group, a substantial number of regenerated plants proved not to have a transgene integrated in their nuclear DNA.

- 36 -

Table 4:

Treatment	Id. No.	Vegetative phenotype ^a	Reproductive phenotype ^b	Copy No. of the transgenes ^c	Phenotype of the F1-progeny ^d
no treatment	1	stressed	sterile	3	stressed/sterile
	2	stressed	sterile	4-6	ND
	3	stressed	sterile	3	stressed/sterile
	4	normal	sterile	1	normal/sterile
	5	stressed	(bud fall)	ND	ND
Before 250	1	normal	sterile	1	normal/sterile
	2	normal	sterile	3	normal/sterile
	3	normal	sterile	1	ND
	4	normal	sterile	3	ND
	5	normal	sterile	2	ND

- a. Vegetatively stressed plants have a small size and flower early, leaves are oblong and dark green.
- b. Reproductive phenotype regards male sterility; in flowers where the buds fell off prematurely this phenotype was not scored, except where some buds resulted in flowers.
- c. Copy number of the transgenes was estimated by comparative Southern. ND: not determined.
- d. F1-progeny was obtained by pollinating the transformed plants with pollen obtained from an untransformed N90-740 line. F1-Progeny resistant to phosphinotricin was scored for vegetative and reproductive phenotype.

- 37 -

Example 4: Agrobacterium-mediated transformation of oilseed rape using niacinamide in the culture medium.

Hypocotyl explants of Brassica napus were obtained as described in Example 3. Four groups of 200 hypocotyl explants each, were either not treated with niacinamide (indicated in table 4 as NONE), treated with 250 mg/l niacinamide for 1 day prior to infection with Agrobacterium (BEFORE), treated for 2 days during the infection with 250 mg/l niacinamide (DURING), or treated for 1 day after the Agrobacterium infection with 250 mg/l niacinamide (AFTER).

All hypocotyl explants were infected with Agrobacterium tumefaciens strain C58C1Rif carrying T-DNA vector pTHW142 and a helper Ti-plasmid pMP90 (Koncz and Shell, 1986 supra)(or a derivative thereof).

Plasmid pTHW142 is a vector carrying a T-DNA comprising the following chimeric genes:

- PSSU-bar-3'g7
- p35S-uidA-3'35S

In which uidA is a DNA encoding b-glucuronidase (Jefferson et al., 1986, Proc. Natl. Acad. Sci. USA 83, 8447-8451) and 3' 35S is the 3' untranslated end of the cauliflower mosaic virus 35S transcript.

The complete sequence of the T-DNA of pTHW142 is presented in SEQ ID No 5.

After the Agrobacterium infection, hypocotyl explants were transferred to selection medium A5, and if appropriate to A5 medium containing 250 mg/l niacinamide. The hypocotyl explants that were placed on medium containing niacinamide were transferred after 1 day to niacinamide-free selection medium A5. After 5 weeks on selective medium the number of transformed calli was scored. b-glucuronidase expression was verified in the obtained calli using established protocols (Jefferson et al., 1986). The results are summarized in

Table 5. Niacinamide treatment either before or after the Agrobacterium infection significantly increase the transformation efficiency.

Table 5:

Treatment	Transformation frequency ^a	Remarks ^b
NONE	16%	small, green calli
BEFORE	32%	large, green calli
DURING	16%	very small, light green calli large, green calli
AFTER	29%	developing shoots

a. Determined as the number of transformed calli (PPT-resitant and GUS-positive) developing per 100 hypocotyl explants

b. Size determination was as follows:

very small: callus diameter of approximately 1-2 mm

small: callus diameter of approximately 2-3 mm

large: callus diameter of approximately 5 mm

All publications cited in this application are hereby incorporated by reference.

- 39 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: PLANT GENETIC SYSTEMS N.V.
- (B) STREET: Plateaustraat 22
- (C) CITY: Ghent
- (E) COUNTRY: Belgium
- (F) POSTAL CODE (ZIP): 9000
- (G) TELEPHONE: 32 9 235 84 58
- (H) TELEFAX: 32 9 223 19 23
- (I) TELEX: 11.361 Pgsgen

(ii) TITLE OF INVENTION: Genetic Transformation using a PARP inhibitor

(iii) NUMBER OF SEQUENCES: 5

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 4946 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: T-DNA of plasmid pTHW107

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION: complement (1..25)
- (D) OTHER INFORMATION: /label= RB
/note= "T-DNA right border"

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION: complement (97..330)
- (D) OTHER INFORMATION: /label= 3'g7
/note= "3' untranslated region containing the
polyadenylation signal of gene 7 of Agrobacterium T-DNA "

- 40 -

(ix) FEATURE:

(A) NAME/KEY: -

(B) LOCATION:complement (331..882)

(D) OTHER INFORMATION:/label= bar

/note= "region coding for phosphinothricin acetyl
transferase"

(ix) FEATURE:

(A) NAME/KEY: -

(B) LOCATION:complement (883..2608)

(D) OTHER INFORMATION:/label= PSSU

/note= "promoter region of Rubisco small subunit gene of
Arabidopsis thali..."

(ix) FEATURE:

(A) NAME/KEY: -

(B) LOCATION:complement (2658..3031)

(D) OTHER INFORMATION:/label= 3'nos

/note= "3' untranslated region containing the
polyadenylation signal of the nopaline synthase gene of Agrobacterium
T-DNA"

(ix) FEATURE:

(A) NAME/KEY: -

(B) LOCATION:complement (3032..3367)

(D) OTHER INFORMATION:/label= barnase

/note= "region coding for barnase"

(ix) FEATURE:

(A) NAME/KEY: -

(B) LOCATION:complement (3368..4876)

(D) OTHER INFORMATION:/label= PTA29

/note= "promoter region of TA29 gene of Nicotiana tabacum"

(ix) FEATURE:

(A) NAME/KEY: -

(B) LOCATION:complement (4922..4946)

(D) OTHER INFORMATION:/label= LB

/note= "T-DNA left border"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

AATTACAACG GTATATATCC TGCCAGTACT CGGCCGTCGA ACTCGGCCGT CGAGTACATG	60
GTCGATAAGA AAAGGCAATT TGTAGATGTT AATTCCCATC TTGAAAGAAA TATAGTTTAA	120
ATATTTATTG ATAAAATAAC AAGTCAGGTA TTATAGTCCA AGCAAAAACA TAAATTTATT	180
GATGCAAGTT TAAATTCAGA AATATTTCAA TAACTGATTA TATCAGCTGG TACATTGCCG	240
TAGATGAAAG ACTGAGTGCG ATATTATGTG TAATACATAA ATTGATGATA TAGCTAGCTT	300
AGCTCATCGG GGGATCCTAG ACGCGTGAGA TCAGATCTCG GTGACGGGCA GGACCGGACG	360

- 41 -

GGGCGGTACC GGCAGGCTGA AGTCCAGCTG CCAGAAACCC ACGTCATGCC AGTTCCTG 420
 CTTGAAGCCG GCCGCCCCGA GCATGCCGCG GGGGGCATAT CCGAGCGCCT CGTGCATGCG 480
 5 CACGCTCGGG TCGTTGGGCA GCCCAGATGAC AGCGACCACG CTCTTGAAGC CCTGTGCCTC 540
 CAGGGACTTC AGCAGGTGGG TGTAGAGCGT GGAGCCAGT CCCGTCCGCT GGTGGCGGGG 600
 10 GGAGACGTAC ACGGTCGACT CGGCCGTCCA GTCGTAGGCG TTGCGTGCCT TCCAGGGGCC 660
 CGCGTAGGCG ATGCCGGCGA CCTCGCCGTC CACCTCGGCG ACGAGCCAGG GATAGCGCTC 720
 CCGCAGACGG ACGAGGTTCGT CCGTCCACTC CTGCGGTTCC TGC GGCTCGG TACGGAAGTT 780
 15 GACCGTGCTT GTCTCGATGT AGTGGTTGAC GATGGTGCAG ACCGCCGCA TGTCCGCCTC 840
 GGTGGCACGG CGGATGTCGG CCGGGCGTCG TTCTGGGTCC ATTGTTCTTC TTTACTCTTT 900
 20 GTGTGACTGA GGT TTGGTCT AGTGCTTTGG TCATCTATAT ATAATGATAA CAACAATGAG 960
 AACAAGCTTT GGAGTGATCG GAGGGTCTAG GATACATGAG ATTCAAGTGG ACTAGGATCT 1020
 ACACCGTTGG ATTTTGAGTG TGGATATGTG TGAGGTTAAT TTTACTTGGT AACGGCCACA 1080
 25 AAGGCCTAAG GAGAGGTGTT GAGACCCTTA TCGGCTTGAA CCGCTGGAAT AATGCCACGT 1140
 GGAAGATAAT TCCATGAATC TTATCGTTAT CTATGAGTGA AATTGTGTGA TGGTGGAGTG 1200
 30 GTGCTTGCTC ATTTTACTTG CCTGGTGGAC TTGGCCCTTT CCTTATGGGG AATTTATATT 1260
 TTTACTTACTA TAGAGCTTTC ATACCTTTTT TTTACCTTGG ATTTAGTTAA TATATAATGG 1320
 TATGATTCAT GAATAAAAAT GGGAAATTTT TGAATTTGTA CTGCTAAATG CATAAGATTA 1380
 35 GGTGAACTG TGGAATATAT ATTTTTTTTCA TTTAAAAGCA AAATTTGCCT TTTACTAGAA 1440
 TTATAAATAT AGAAAAATAT ATAACATTCA AATAAAAATG AAAATAAGAA CTTTCAAAAA 1500
 40 ACAGAACTAT GTTTAATGTG TAAAGATTAG TCGCACATCA AGTCATCTGT TACAATATGT 1560
 TACAACAAGT CATAAGCCCA ACAAGTTAG CACGTCTAAA TAACTAAAG AGTCCACGAA 1620
 AATATTACAA ATCATAAGCC CAACAAAGTT ATTGATCAAA AAAAAAAAC GCCCAACAAA 1680
 45 GCTAAACAAA GTCCAAAAAA AACTTCTCAA GTCTCCATCT TCCTTTATGA ACATTGAAAA 1740
 CTATACACAA AACAAGTCAG ATAAATCTCT TTCTGGGCCT GTCTTCCCA CCTCCTACAT 1800
 50 CACTTCCCTA TCGGATTGAA TGTTTTACTT GTACCTTTTC CGTTGCAATG ATATTGATAG 1860
 TATGTTTGTG AAAACTAATA GGGTTAACAA TCGAAGTCAT GGAATATGGA TTTGGTCCAA 1920
 55 GATTTTCCGA GAGCTTTCTA GTAGAAAGCC CATCACCAGA AATTTACTAG TAAAATAAAT 1980

- 42 -

	CACCAATTAG	GTTTCTTATT	ATGTGCCAAA	TTCAATATAA	TTATAGAGGA	TATTTCAAAT	2040
	GAAAACGTAT	GAATGTTATT	AGTAAATGGT	CAGGTAAGAC	ATTAAAAAAA	TCCTACGTCA	2100
5	GATATTCAAC	TTTAAAAATT	CGATCAGTGT	GGAATTGTAC	AAAAATTTGG	GATCTACTAT	2160
	ATATATATAA	TGCTTTACAA	CAC TTGGATT	TTTTTTTGGG	GGCTGGAATT	TTTAATCTAC	2220
	ATATTTGTTT	TGGCCATGCA	CCAAC TCATT	GTTTAGTGTA	ATACTTTGAT	TTTGTC AAAT	2280
10	ATATGTGTTT	GTGTATATTT	GTATAAGAAT	TTCTTTGACC	ATATACACAC	ACACATATAT	2340
	ATATATATAT	ATATATTATA	TATCATGCAC	TTTTAATTGA	AAAAATAATA	TATATATATA	2400
15	TAGTGCATTT	TTTCTAACAA	CCATATATGT	TGCGATTGAT	CTGCAAAAAT	ACTGCTAGAG	2460
	TAATGAAAAA	TATAATCTAT	TGCTGAAATT	ATCTCAGATG	TTAAGATTTT	CTTAAAGTAA	2520
	ATTCTTTCAA	ATTTTAGCTA	AAAGTCTTGT	AATAACTAAA	GAATAATACA	CAATCTCGAC	2580
20	CACGGAAAAA	AAACACATAA	TAAATTTGAA	TTTCGACCGC	GGTACCCGGA	ATTCGAGCTC	2640
	GGTACCCGGG	GATCTTCCCG	ATCTAGTAAC	ATAGATGACA	CCGCGCGCGA	TAATTTATCC	2700
25	TAGTTTGCGC	GCTATATTTT	GTTTTCTATC	GCGTATTAAA	TGTATAATTG	CGGGACTCTA	2760
	ATCATAAAAA	CCCATCTCAT	AAATAACGTC	ATGCATTACA	TGTTAATTAT	TACATGCTTA	2820
	ACGTAATTCA	ACAGAAATTA	TATGATAATC	ATCGCAAGAC	CGGCAACAGG	ATTCAATCTT	2880
30	AAGAACTTTT	ATTGCCAAAT	GTTTGAACGA	TCTGCTTCGG	ATCCTCTAGA	GCCGGAAAGT	2940
	GAAATTGACC	GATCAGAGTT	TGAAGAAAAA	TTTATTACAC	ACTTTATGTA	AAGCTGAAAA	3000
35	AAACGGCCTC	CGCAGGAAGC	CGTTTTTTTC	GTTATCTGAT	TTTTGTAAAG	GTCTGATAAT	3060
	GGTCCGTTGT	TTTGTAATC	AGCCAGTCGC	TTGAGTAAAG	AATCCGGTCT	GAATTTCTGA	3120
	AGCCTGATGT	ATAGTTAATA	TCCGCTTCAC	GCCATGTTCT	TCCGCTTTTG	CCCGGGAGTT	3180
40	TGCCTTCCCT	GTTTGAGAAG	ATGTCTCCGC	CGATGCTTTT	CCCCGGAGCG	ACGTCTGCAA	3240
	GGTTCCTTTT	TGATGCCACC	CAGCCGAGGG	CTTG TGCTTC	TGATTTTGTA	ATGTAATTAT	3300
45	CAGGTAGCTT	ATGATATGTC	TGAAGATAAT	CCGCAACCCC	GTCAAACGTG	TTGATAACCG	3360
	GTACCATGGT	AGCTAATTTT	TTTAAGTAAA	AAC TTGATT	TGAGTGATGA	TGTTG TACTG	3420
	TTACACTTGC	ACCACAAGGG	CATATATAGA	GCACAAGACA	TACACAACAA	CTTGCAAAAC	3480
50	TAAC TTTTGT	TGGAGCATTT	CGAGGAAAAT	GGGGAGTAGC	AGGCTAATCT	GAGGGTAACA	3540
	TTAAGGTTTC	ATGTATTAAT	TTGTTGCAAA	CATGGACTTA	GTGTGAGGAA	AAAGTACCAA	3600
55	AATTTTGTCT	CACCCTGATT	TCAGTTATGG	AAATTACATT	ATGAAGCTGT	GCTAGAGAAG	3660

- 43 -

ATGTTTATTC TAGTCCAGCC ACCCACCTTA TGCAAGTCTG CTTTGTAGCTT GATTCAAAAA 3720
CTGATTTAAT TTACATTGCT AAATGTGCAT ACTTCGAGCC TATGTCGCTT TAATTCGAGT 3780
5 AGGATGTATA TATTAGTACA TAAAAAATCA TGTTTGAATC ATCTTTCATA AAGTGACAAG 3840
TCAATTGTCC CTTCTTGTTT GGCACATAT TCAATCTGTT AATGCAAATT ATCCAGTTAT 3900
10 ACTTAGCTAG ATATCCAATT TTGAATAAAA ATAGCTCTTG ATTAGTAAAC CGGATAGTGA 3960
CAAAGTCACA TATCCATCAA ACTTCTGGTG CTCGTGGCTA AGTTCTGATC GACATGGGGT 4020
TAAAATTTAA ATTGGGACAC ATAAATAGCC TATTTGTGCA AATCTCCCCA TCGAAAATGA 4080
15 CAGATTGTTA CATGGAAAAC AAAAAGTCCT CTGATAGAAG TCGCAAAGTA TCACAATTTT 4140
CTATCGAGAG ATAGATTGAA AGAAGTGCAG GGAAGCGGTT AACTGGAACA TAACACAATG 4200
20 TCTAAATTAA TTGCATTGCG TAACCAAAAA GTGTATTACT CTCTCCGGTC CACAATAAGT 4260
TATTTTTTGG CCCTTTTTTT ATGGTCCAAA ATAAGTGAGT TTTTGTAGATT TCAAAAATGA 4320
TTTAATTATT TTTTACTAC AGTGCCCTTG GAGTAAATGG TGTGGAGTA TGTGTTAGAA 4380
25 ATGTTTATGT GAAGAAATAG TAAAGGTAA TATGATCAAT TTCATTGCTA TTTAATGTTA 4440
AAATGTGAAT TTCTTAATCT GTGTGAAAAC AACCAAAAAA TCACTTATTG TGGACCGGAG 4500
30 AAAGTATATA AATATATATT TGGAAGCGAC TAAAAATAAA CTTTCTCAT ATTATACGAA 4560
CCTAAAAACA GCATATGGTA GTTCTAGGG AATCTAAATC ACTAAAATTA ATAAAAGAAG 4620
CAACAAGTAT CAATACATAT GATTTACACC GTCAAACACG AAATTCGTAA ATATTTAATA 4680
35 TAATAAGAA TTAATCCAAA TAGCCTCCCA CCCTATAACT TAAACTAAAA ATAACCAGCG 4740
AATGTATATT ATATGCATAA TTTATATATT AAATGTGTAT AATCATGTAT AATCAATGTA 4800
40 TAATCTATGT ATATGGTTAG AAAAAGTAAA CAATTAATAT AGCCGGCTAT TTGTGTAAAA 4860
ATCCCTAATA TAATCGCGAC GGATCCCCGG GAATTCCGGG GAAGCTTAGA TCCATGGAGC 4920
CATTTACAAT TGAATATATC CTGCCG 4946

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6548 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: circular

(ii) MOLECULE TYPE: DNA (genomic)

- 44 -

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

5 (vi) ORIGINAL SOURCE:
(A) ORGANISM: plasmid pTS172

(ix) FEATURE:
10 (A) NAME/KEY: -
(B) LOCATION:complement (2019..2288)
(D) OTHER INFORMATION:/label= 3'nos
/note= "3' untranslated region containing the
polyadenylation signal of the nopaline synthase gene of Agrobacterium
T-DNA"

15 (ix) FEATURE:
(A) NAME/KEY: -
(B) LOCATION:complement (2289..2624)
(D) OTHER INFORMATION:/label= barnase
20 /note= "region coding for barnase"

(ix) FEATURE:
(A) NAME/KEY: -
(B) LOCATION:complement (2625..4313)
25 (D) OTHER INFORMATION:/label= PE1
/note= "promoter region of E1 gene of rice"

(ix) FEATURE:
(A) NAME/KEY: -
(B) LOCATION:4336..5710
30 (D) OTHER INFORMATION:/label= P35S
/note= "35S promoter region of Cauliflower mosaic virus"

(ix) FEATURE:
35 (A) NAME/KEY: -
(B) LOCATION:5711..6262
(D) OTHER INFORMATION:/label= bar
/note= "region coding for phosphinothricin acetyl
40 transferase"

(ix) FEATURE:
(A) NAME/KEY: -
(B) LOCATION:6263..6496
45 (D) OTHER INFORMATION:/label= 3'g7
/note= "3' untranslated region containing the
polyadenylation signal of gene 7 of Agrobacterium T-DNA"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

50 AATTCAAGCT TGACGTCAGG TGGCACTTTT CGGGGAAATG TGC GCGGAAC CCCTATTTGT 60
TTATTTTCT AAATACATTC AAATATGTAT CCGCTCATGA GACAATAACC CTGATAAATG 120
55 CTTCAATAAT ATTGAAAAAG GAAGAGTATG AGTATTCAAC ATTTCCGTGT CGCCCTTATT 180

- 45 -

CCCTTTTTTG CGGCATTTTG CCTTCCTGTT TTTGCTCACC CAGAAACGCT GGTGAAAGTA 240
AAAGATGCTG AAGATCAGTT GGGTGCACGA GTGGGTTACA TCGAACTGGA TCTCAACAGC 300
5 GGTAAGATCC TTGAGAGTTT TCGCCCCGAA GAACGTTTTT CAATGATGAG CACTTTTAAA 360
GTTCTGCTAT GTGGCGCGGT ATTATCCCGT ATTGACGCCG GGCAAGAGCA ACTCGGTTCG 420
10 CGCATACACT ATTCTCAGAA TGACTTGGTT GAGTACTCAC CAGTCACAGA AAAGCATCTT 480
ACGGATGGCA TGACAGTAAG AGAATTATGC AGTGCTGCCA TAACCATGAG TGATAACACT 540
GCGGCCAACT TACTTCTGAC AACGATCGGA GGACCGAAGG AGCTAACCGC TTTTTTGCAC 600
15 AACATGGGGG ATCATGTAAC TCGCCTTGAT CGTTGGGAAC CGGAGCTGAA TGAAGCCATA 660
CCAAACGACG AGCGTGACAC CACGATGCCT GTAGCAATGG CAACAACGTT GCGCAAACATA 720
TTAACTGGCG AACTACTTAC TCTAGCTTCC CGGCAACAAT TAATAGACTG GATGGAGGCG 780
20 GATAAAGTTG CAGGACCACT TCTGCGCTCG GCCCTTCCGG CTGGCTGGTT TATTGCTGAT 840
AAATCTGGAG CCGGTGAGCG TGGGTCTCGC GGTATCATTG CAGCACTGGG GCCAGATGGT 900
25 AAGCCCTCCC GTATCGTAGT TATCTACACG ACGGGGAGTC AGGCAACTAT GGATGAACGA 960
AATAGACAGA TCGCTGAGAT AGGTGCCTCA CTGATTAAGC ATTGGTAACT GTCAGACCAA 1020
30 GTTTACTCAT ATATACTTTA GATTGATTTA AAACCTTCATT TTTAATTTAA AAGGATCTAG 1080
GTGAAGATCC TTTTGGGCTC GAGTCTCATG ACCAAAATCC CTTAACGTGA GTTTTCGTTC 1140
CACTGAGCGT CAGACCCCGT AGAAAAGATC AAAGGATCTT CTTGAGATCC TTTTTTCTG 1200
35 CGCGTAATCT GCTGCTTGCA AACAAAAAAA CCACCGCTAC CAGCGGTGGT TTGTTTGCCG 1260
GATCAAGAGC TACCAACTCT TTTCCGAAG GTAAGTGGCT TCAGCAGAGC GCAGATACCA 1320
40 AATACTGTCC TTCTAGTGTA GCCGTAGTTA GGCCACCACT TCAAGAACTC TGTAGCACCG 1380
CCTACATACC TCGCTCTGCT AATCCTGTTA CCAGTGCGTG CTGCCAGTGG CGATAAGTCG 1440
TGTCTTACCG GGTGGAATC AAGACGATAG TTACCGGATA AGGCGCAGCG GTCGGGCTGA 1500
45 ACGGGGGGTT CGTGACACA GCCCAGCTTG GAGCGAACGA CCTACACCGA ACTGAGATAC 1560
CTACAGCGTG AGCATTGAGA AAGCGCCACG CTTCCCGAAG GGAGAAAGGC GGACAGGTAT 1620
50 CCGGTAAGCG GCAGGGTCGG AACAGGAGAG CGCACGAGGG AGCTTCCAGG GGGAAACGCC 1680
TGGTATCTTT ATAGTCCTGT CGGGTTTCGC CACCTCTGAC TTGAGCGTCG ATTTTTGTGA 1740
TGCTCGTCAG GGGGGCGGAG CCTATGGAAA AACGCCAGCA ACGCGGCCTT TTTACGGTTC 1800
55

- 46 -

	CTGGCCTTTT	GCTGGCCTTT	TGCTCACATG	TTCTTTCCTG	CGTTATCCCC	TGATTCTGTG	1860
	GATAACCGTA	TTACCGCCTT	TGAGTGAGCT	GATACCGCTC	GCCGCAGCCG	AACGACCGAG	1920
5	CGCAGCGAGT	CAGTGAGCGA	GGAAGCGGAA	GAGCGCCCAA	TACGCAAACC	GCCTCTCCCC	1980
	GCGCGTTGGC	CTGATCAGAA	TTCATATGCA	CGTGTTCCTG	ATCTAGTAAC	ATAGATGACA	2040
	CCGCGCGCGA	TAATTTATCC	TAGTTTGCGC	GCTATATTTT	GTTTTCTATC	GCGTATTAAA	2100
10	TGTATAATTG	CGGGACTCTA	ATCATAAAAA	CCCATCTCAT	AAATAACGTC	ATGCATTACA	2160
	TGTTAATTAT	TACATGCTTA	ACGTAATTCA	ACAGAAATTA	TATGATAATC	ATCGCAAGAC	2220
15	CGGCAACAGG	ATTCAATCTT	AAGAACTTTT	ATTGCCAAAT	GTTTGAACGA	TCTGCTTCGG	2280
	AGGTTACCTT	ATCTGATTTT	TGTAAAGGTC	TGATAATGGT	CCGTTGTTTT	GTAAATCAGC	2340
	CAGTCGCTTG	AGTAAAGAAT	CCGGTCTGAA	TTTCTGAAGC	CTGATGTATA	GTAAATATCC	2400
20	GCTTCACGCC	ATGTTCTGTC	GCTTTTGCCC	GGGAGTTTGC	CTTCCCTGTT	TGAGAAGATG	2460
	TCTCCGCCGA	TGCTTTTCCC	CGGAGCGACG	TCTGCAAGGT	TCCCTTTTGA	TGCCACCCAG	2520
25	CCGAGGGCTT	GTGCTTCTGA	TTTTGTAATG	TAATTATCAG	GTAGCTTATG	ATATGTCTGA	2580
	AGATAATCCG	CAACCCCGTC	AAACGTGTTG	ATAACCGGTA	CCATCGCGAC	GGCTTGATGG	2640
	ATCTCTTGCT	GGACACCGGG	ATGCTAGGAT	GGGTTATCGT	GGCCGGCGTG	CGTGTGTGGC	2700
30	TTTTGTAGGC	GCCGGCGACG	GCGGGGGCAA	TGTGGCAGGT	GAGTCACGGT	GCAAGCGTGC	2760
	GCAAGTGAAT	GCAACAACCA	AGGACGGTCA	TGGCGAAAGC	ACCTCACGCG	TCCACCGTCT	2820
35	ACAGGATGTA	GCAGTAGCAC	GGTGAAAGAA	GTGTTGTCCC	GTCCATTAGG	TGCATTCTCA	2880
	CCGTTGGCCA	GAACAGGACC	GTTCAACAGT	TAGGTTGAGT	GTAGGACTTT	TACGTGGTTA	2940
	ATGTATGGCA	AATAGTAGTA	AATTTTGCCC	CCATTGGTCT	GGCTGAGATA	GAACATATTC	3000
40	TGGAAAGCCT	CTAGCATATC	TTTTTTGACA	GCTAAACTTT	GCTTCTTGCC	TTCTTGGTCT	3060
	AGCAATGACG	TTGCCCATGT	CGTGGCAAAC	ATCTGGTAAG	GTAAGTGTAT	TCGTTTGTTC	3120
45	CCTTCAACGG	CTCAATCCCC	ACAGGCCAAG	CTATCCTTTC	CTTGGCAGTA	TAGGCTCCTT	3180
	GAGAGATTAT	ACTACCATTT	TTAAGTGCTT	ATAAAGACGA	TGCTCTCTAA	CCAGATCGAT	3240
	CAGAAACACA	AAGTTTTAGC	AGCGTAATAT	CCCACACACA	TACACACACG	AAGCTATGCC	3300
50	TCCTCATTTT	CCGAGAGATT	CTGACAGTGA	CCAGAATGTC	AGAATGCCAT	TTCATGGGCA	3360
	CAAGTCGATC	CACAAGCTTC	TTGGTGGAGG	TCAAGGTGTG	CTATTATTAT	TCGCTTTCTA	3420
55	GGAAATTATT	CAGAATTAGT	GCCTTTTATC	ATAACTTCTC	TCTGAGCCGA	TGTGGTTTTG	3480

- 47 -

	GATTTTCATTG TTGGGAGCTA TGCAGTTGCG GATATTCTGC TGTGGAAGAA CAGGAACTTA	3540
	TCTGCGGGGG TCCTTGCTGG GGCAACATTG ATATGGTTCC TGTTGATGT AGTAGAATAC	3600
5	AATATAATTC CGCTCCTTTG CCAGATTGCC ATTCTTGCCA TGCTTGTGAT CTTCAATTTGG	3660
	TCAAATGCCG CACCACTCTT GGACAGGTAT TAGCTTTATT TCCTGTGGAG ATGGTAGAAA	3720
10	ACTCAGCTTA CAGAAATGGC ATTTACGTA GTATAACGCA AGACATTAGG TACTAAACT	3780
	CAACTAACTG TTTCCGAATT TCAGGGCCCC TCCAAGGATC CCAGAAATCA TCATCTCTGA	3840
	ACATGCCTTC AGAGAAATGG CATTGACCGT CCATTACAAA CTAACGTACA CTGTATCTGT	3900
15	TCTTTACGAC ATTGCATGTG GAAAGGATCT GAAGAGATTT CTCCTGGTAC ATAATAATCT	3960
	ACTCCTTTGC TACGTTAATA AGAGATGTAA AAACATGCAA CAGTTCCAGT GCCAACATTG	4020
20	TCCAAGGATT GTGCAATTCT TTCTGGAGCG CTAAAATTGA CCAGATTAGA CGCATCAGAA	4080
	TATTGAATTG CAGAGTTAGC CAATAATCCT CATAATGTTA ATGTGCTATT GTTGTTCACT	4140
	ACTCAATATA GTTCTGGACT AACAAATCAGA TTGTTTATGA TATTAAGGTG GTTGGATCTC	4200
25	TATTGGTATT GTCGGCGATT GGAAGTTCTT GCAGCTTGAC AAGTCTACTA TATATTGGTA	4260
	GGTATTCCAG ATAAATATTA AATTTTAATA AAACAATCAC ACAGAAGGAT CTGCGGCCGC	4320
30	TAGCCTAGGC CCGGGCCAC AAAAATCTGA GCTTAACAGC ACAGTTGCTC CTCTCAGAGC	4380
	AGAATCGGGT ATTCAACACC CTCATATCAA CTAATACGTT GTGTATAACG GTCCACATGC	4440
	CGGTATATAC GATGACTGGG GTTGATACAAA GGCGGCAACA AACGGCGTTC CCGGAGTTGC	4500
35	ACACAAGAAA TTTGCCACTA TTACAGAGGC AAGAGCAGCA GCTGACGCGT ACACAACAAG	4560
	TCAGCAAACA GACAGGTTGA ACTTCATCCC CAAAGGAGAA GCTCAACTCA AGCCCAAGAG	4620
40	CTTTGCTAAG GCCCTAACAA GCCCACCAAA GCAAAAAGCC CACTGGCTCA CGCTAGGAAC	4680
	CAAAAGGCCC AGCAGTGATC CAGCCCCAAA AGAGATCTCC TTTGCCCCGG AGATTACAAT	4740
45	GGACGATTC CTCTATCTTT ACGATCTAGG AAGGAAGTTC GAAGGTGAAG GTGACGACAC	4800
	TATGTTCACT ACTGATAATG AGAAGGTTAG CCTCTTCAAT TTCAGAAAGA ATGCTGACCC	4860
	ACAGATGGTT AGAGAGGCCT ACGCAGCAGG TCTCATCAAG ACGATCTACC CGAGTAACAA	4920
50	TCTCCAGGAG ATCAAATACC TTCCAAGAA GGTAAAGAT GCAGTCAAAA GATTCAGGAC	4980
	TAATTGCATC AAGAACACAG AGAAAGACAT ATTTCTCAAG ATCAGAAGTA CTATTCCAGT	5040
55	ATGGACGATT CAAGGCTTGC TTCATAAACC AAGGCAAGTA ATAGAGATTG GAGTCTCTAA	5100

- 48 -

AAAGGTAGTT CCTACTGAAT CTAAGGCCAT GCATGGAGTC TAAGATTCAA ATCGAGGATC 5160
TAACAGAACT CGCCGTGAAG ACTGGCGAAC AGTTCATACA GAGTCTTTTA CGACTCAATG 5220
5 ACAAGAAGAA AATCTTCGTC AACATGGTGG AGCACGACAC TCTGGTCTAC TCCAAAAATG 5280
TCAAAGATAC AGTCTCAGAA GACCAAAGGG CTATTGAGAC TTTTCAACAA AGGATAATTT 5340
CGGGAAACCT CCTCGGATTC CATTGCCCAG CTATCTGTCA CTTTCATCGAA AGGACAGTAG 5400
10 AAAAGGAAGG TGGCTCCTAC AAATGCCATC ATTGCGATAA AGGAAAGGCT ATCATTCAAG 5460
ATGCCTCTGC CGACAGTGGT CCCAAAGATG GACCCCCACC CACGAGGAGC ATCGTGAAAA 5520
15 AAGAAGACGT TCCAACCACG TCTTCAAAGC AAGTGGATTG ATGTGACATC TCCACTGACG 5580
TAAGGGATGA CGCACAATCC CACTATCCTT CGCAAGACCC TTCCTCTATA TAAGGAAGTT 5640
CATTTCAATTT GGAGAGGACA CGCTGAAATC ACCAGTCTCT CTCTATAAAT CTATCTCTCT 5700
20 CTCTATAACC ATGGACCCAG AACGACGCCC GGCCGACATC CGCCGTGCCA CCGAGGCGGA 5760
CATGCCGGCG GTCTGCACCA TCGTCAACCA CTACATCGAG ACAAGCACGG TCAACTTCCG 5820
25 TACCGAGCCG CAGGAACCGC AGGAGTGGAC GGACGACCTC GTCCGTCTGC GGGAGCGCTA 5880
TCCCTGGCTC GTCGCCGAGG TGGACGGCGA GGTCGCCGGC ATCGCCTACG CGGGCCCCCTG 5940
GAAGGCACGC AACGCCTACG ACTGGACGGC CGAGTCGACC GTGTACGTCT CCCCCGCCA 6000
30 CCAGCGGACG GGAAGTGGCT CCACGCTCTA CACCCACCTG CTGAAGTCCC TGGAGGCACA 6060
GGGCTTCAAG AGCGTGGTCG CTGTCATCGG GCTGCCCAAC GACCCGAGCG TGCATGCA 6120
35 CGAGGCGCTC GGATATGCCC CCCGCGGCAT GCTGCGGGCG GCCGGCTTCA AGCACGGGAA 6180
CTGGCATGAC GTGGGTTTCT GGCAGCTGGA CTTACAGCTG CCGGTACCGC CCCGTCCGGT 6240
CCTGCCCCGTC ACCGAGATCT GAGATCACGC GTTCTAGGAT CCCCCGATGA GCTAAGCTAG 6300
40 CTATATCATC AATTTATGTA TTACACATAA TATCGCACTC AGTCTTTCAT CTACGGCAAT 6360
GTACCAGCTG ATATAATCAG TTATTGAAAT ATTTCTGAAT TTAAACTTGC ATCAATAAAT 6420
45 TTATGTTTTT GCTTGGACTA TAATACCTGA CTTGTTATTT TATCAATAAA TATTTAAACT 6480
ATATTTCTTT CAAGATGGGA ATTAACATCT ACAAATTGCC TTTTCTTATC GACCATGTAC 6540
GTATCGCG 6548

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1601 base pairs

(B) TYPE: nucleic acid

- 49 -

(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: T72 promoter region

(ix) FEATURE:

(A) NAME/KEY: -

(B) LOCATION: complement (1..1601)

(D) OTHER INFORMATION: /label= PT72

/note= "promoter region of T72 gene of rice"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

CGCCGTGAGT	GTCTTCTGCC	GCCGAGGGGC	TCTCGCTCGT	CGTCGATGCC	TGCACGGTGC	60
GTGCGTGTGT	GTCGTGGTGG	TGGTGGCGAT	ACGCGACGCG	AGCTCGATTT	ATAGGAGGGG	120
ATCGAAGGAG	GGGAGCGCGC	GCGGCGAGGC	CCGCGTTGCT	CACCTACGCC	GCGCGCATGC	180
GGCGGACGCG	CGGTCGGCGC	CCGCGCCGGC	CGGGAGGACG	AGGGCGCAAG	CGTGTGAGCC	240
ACCGAACGCG	CGCGCGCGCC	GCGGCGCGAA	CTCTCCATCG	CGTCGCGGCG	AGCCGAGAGC	300
CGACGAGAGC	GTTTCGCGCG	CGCGGTTGGG	CCGGCGACAA	GATGGGCCGT	AGCCCTGGGC	360
CTCGTGCCAT	CTTTTTTTTT	CTTTTTTGCC	TTTTTTGGCC	TGGCAATTTT	TTTTTGTTTT	420
TAGTCTTTTT	GTGGTGATAA	TGTGTCGTCT	TCCGGTGAAC	TAATTTACTC	GTTGATCTTT	480
TTGTGTCCCT	TCGAATATTC	GCAGTGGTAG	AAGATGACTA	CTACTACCAG	TAGTTGATCT	540
CGAATGGCAA	CTTTTGTGCA	GAAC TTATTC	CACGGCTATG	TCAGCTTCCA	CTGTGACTAA	600
AAAAACTACG	GCCATCTTTT	GGA CTGTTC	TATCTTGGA	CTGAACAAAA	AGGACGATCC	660
TGATGTACAC	ACGGCATAGT	TTCCAGCACT	GGATGCCAAG	TTGCCAACTG	TTACCACGAT	720
AATGGAACGA	CGAGATGAGA	TATTATACAA	GTCCAATGGA	TCAAGATCCT	GTGCAGTTGT	780
TATTGTA ACT	GTA ACTTAAG	CCGT TAACAT	GTACATCACA	TTTCCTACTC	TATCAATGTC	840
TTGTGCGGGT	TGTTTCAAAA	AAACATGTAC	ATCACATGAT	CTAGAACGGA	AGGCCAGGAT	900
ATGAAGTGGT	ACTGCAGCAA	AAACACTGTA	GCAGAGATGT	ACTATTATGC	ATGTACTGTA	960
GCAGTCATCT	AGAGCCGTTG	GATCTGAAAA	CGAATGGACA	TGATTGTGTG	CAGTTGCTAT	1020

- 50 -

TGTGCAGTTA CAATAGCAAC TGCATTTGAT CTTAATCCAA GTCCAATACA TGCAGAACAG 1080
 TAGCTACGAG CTGGAAAGGA TGCAAATCTG GGTGACACTG ACAGCAACCG TGGAAGAACA 1140
 5 ACAGCAGCAA AGTCCCAGAG GGATGGCAAT TTGAAGGAAT TTAAATACTC TAATATTACT 1200
 CCACCCGTTA AAAAAACAA CTTGCTACGC ATAATATATG TTCGGATTTA TAGCGAGAAG 1260
 10 TTAATTTTTC ATGAGAAGAA GAATATATAT GTAATATGTA CTAGGAGAGT ACTCGCTTCA 1320
 TAAATATAAA TATTCATAAG TTGTCCAGTG AAGATAGCTT TAGAAAAAAC TAGTTATTTT 1380
 ATTTGTCAAA TTTTAAATTT TGAAGTAGTT AGATTATCTT TCTAGTAGTT CTGATTGGTT 1440
 15 GAAAATGTTT AGATTTTCAT GTGTTAAGAG TTCCGTATCC TAAAAATAGT AATATAATTT 1500
 TAAATCATAT ATATATATAT ATATATATAT ATATATATAT ATATATATAT ATATATATAT 1560
 TGTGAACGG TTTGTGCTCT GGTGCTATC CTGTTCTGTG G 1601

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6291 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: circular

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: plasmid pVE136

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION: complement (425..687)
- (D) OTHER INFORMATION: /label= 3'nos

/note= "3'untranslated region containing the
 polyadenylation signal of the nopaline synthase gene of Agrobacterium
 T-DNA"

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION: complement (803..1138)
- (D) OTHER INFORMATION: /label= barnase
- /note= "region coding for barnase"

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION: complement (1138..2317)
- (D) OTHER INFORMATION: /label= PCa55

- 51 -

/note= "stamen-specific promoter from corn gene CA55"

(ix) FEATURE:

(A) NAME/KEY: -
(B) LOCATION:2355..3187
(D) OTHER INFORMATION:/label= P35S
/note= "35S promoter region of Cauliflower mosaic virus"

(ix) FEATURE:

(A) NAME/KEY: -
(B) LOCATION:3188..3739
(D) OTHER INFORMATION:/label= bar
/note= "region coding for phosphotransferase"

(ix) FEATURE:

(A) NAME/KEY: -
(B) LOCATION:3757..4017
(D) OTHER INFORMATION:/label= 3'nos
/note= "3' untranslated region containing the
polyadenylation
signal of the nopaline synthase gene of Agrobacterium
T-DNA"

(ix) FEATURE:

(A) NAME/KEY: -
(B) LOCATION:699..702
(D) OTHER INFORMATION:/note= "region with unknown
sequence (may contain up to 15 nucleotides)"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

TCGCGCGTTT	CGGTGATGAC	GGTGAAAACC	TCTGACACAT	GCAGCTCCCG	GAGACGGTCA	60
CAGCTTGTCT	GTAAGCGGAT	GCCGGGAGCA	GACAAGCCCG	TCAGGGCGCG	TCAGCGGGTG	120
TTGGCGGGTG	TCGGGGCTGG	CTTAACATG	CGGCATCAGA	GCAGATTGTA	CTGAGAGTGC	180
ACCATATGCG	GTGTGAAATA	CCGCACAGAT	GCGTAAGGAG	AAAATACCGC	ATCAGGCGCC	240
ATTCGCCATT	CAGGCTGCGC	AACTGTTGGG	AAGGGCGATC	GGTGCGGGCC	TCTTCGCTAT	300
TACGCCAGCT	GGCGAAAGGG	GGATGTGCTG	CAAGGCGATT	AAGTTGGGTA	ACGCCAGGGT	360
TTTCCCAGTC	ACGACGTTGT	AAAACGACGG	CCAGTGAATT	CGAGCTCGGT	ACCCGGGGAT	420
CTTCCCGATC	TAGTAACATA	GATGACACCG	CGCGCGATAA	TTTATCCTAG	TTTGCGCGCT	480
ATATTTTGTT	TTCTATCGCG	TATTAAATGT	ATAATTGCGG	GACTCTAATC	ATAAAAACCC	540
ATCTCATAAA	TAACGTCATG	CATTACATGT	TAATTATTAC	ATGCTTAACG	TAATTCAACA	600
GAAATTATAT	GATAATCATC	GCAAGACCGG	CAACAGGATT	CAATCTTAAG	AAACTTTATT	660

- 52 -

	GCCAAATGTT TGAACGATCT GCTTCGGATC CTCTAGAGNN NNCCGGAAAG TGAAATTGAC	720
	CGATCAGAGT TTGAAGAAAA ATTTATTACA CACTTTATGT AAAGCTGAAA AAAACGGCCT	780
5	CCGCAGGAAG CCGTTTTTTT CGTTATCTGA TTTTGTAAA GGTCTGATAA TGGTCCGTTG	840
	TTTTGTAAAT CAGCCAGTCG CTTGAGTAAA GAATCCGGTC TGAATTTCTG AAGCCTGATG	900
10	TATAGTTAAT ATCCGCTTCA CGCCATGTTT GTCCGCTTTT GCCCGGGAGT TTGCCTTCCC	960
	TGTTTGAGAA GATGTCTCCG CCGATGCTTT TCCCCGGAGC GACGTCTGCA AGGTTCCCTT	1020
	TTGATGCCAC CCAGCCGAGG GCTTGTGCTT CTGATTTTGT AATGTAATTA TCAGGTAGCT	1080
15	TATGATATGT CTGAAGATAA TCCGCAACCC CGTCAAACGT GTTGATAACC GGTACCATGG	1140
	CTGCAGCTAG TTAGCTCGAT GTATCTTCTG TATATGCAGT GCAGCTTCTG CGTTTTGGCT	1200
20	GCTTTGAGCT GTGAAATCTC GCTTTCCAGT CCCTGCGTGT TTTATAGTGC TGTACGTTCTG	1260
	TGATCGTGAG CAAACAGGGC GTGCCTCAAC TACTGGTTTG GTTGGGTGAC AGGCGCCAAC	1320
	TACGTGCTCG TAACCGATCG AGTGAGCGTA ATGCAACATT TTTTCTTCTT CTCTCGCATT	1380
25	GGTTTCATCC AGCCAGGAGA CCCGAATCGA ATTGAAATCA CAAATCTGAG GTACAGTATT	1440
	TTTACAGTAC CGTTCGTTCTG AAGGTCTTCG ACAGGTCAAG GTAACAAAAT CAGTTTTAAA	1500
30	TTGTTGTTTC AGATCAAAGA AAATTGAGAT GATCTGAAGG ACTTGGACCT TCGTCCAATG	1560
	AAACACTTGG ACTAATTAGA GGTGAATTGA AAGCAAGCAG ATGCAACCGA AGGTGGTGAA	1620
	AGTGGAGTTT CAGCATTGAC GACGAAAACC TTCGAACGGT ATAAAAAGA AGCCGCAATT	1680
35	AAACGAAGAT TTGCCAAAAA GATGCATCAA CCAAGGGAAG ACGTGCATAC ATGTTTGATG	1740
	AAAACCTCGTA AAAACTGAAG TACGATTCCC CATTCCCCTC CTTTTCTCGT TTCTTTTAAC	1800
40	TGAAGCAAAG AATTTGTATG TATTCCCTCC ATTCCATATT CTAGGAGGTT TTGGCTTTTC	1860
	ATACCCTCCT CCATTTCAAA TTATTTGTCA TACATTGAAG ATATACACCA TTCTAATTTA	1920
	TACTAAATTA CAGCTTTTAG ATACATATAT TTTATTATAC ACTTAGATAC GTATTATATA	1980
45	AAACACCTAA TTTAAAATAA AAAATTATAT AAAAAGTGTA TCTAAAAAAT CAAAATACGA	2040
	CATAATTTGA AACGGAGGGG TACTACTTAT GCAAACCAAT CGTGGTAACC CTAAACCCTA	2100
50	TATGAATGAG GCCATGATTG TAATGCACCG TCTGATTAAC CAAGATATCA ATGGTCAAAG	2160
	ATATACATGA TACATCCAAG TCACAGCGAA GGCAAATGTG ACAACAGTTT TTTTACCAG	2220
	AGGGACAAGG GAGAATATCT ATTCAGATGT CAAGTCCCG TATCACACTG CCAGGTCCTT	2280
55	ACTCCAGACC ATCTTCCGGC TCTATTGATG CATACCAGGA ATTGATCTAG AGTCGACCTG	2340

- 53 -

	CAGGCATGCA	AGCTCCTACG	CAGCAGGTCT	CATCAAGACG	ATCTACCCGA	GTAACAATCT	2400
	CCAGGAGATC	AAATACCTTC	CCAAGAAGGT	TAAAGATGCA	GTCAAAAGAT	TCAGGACTAA	2460
5	TTGCATCAAG	AACACAGAGA	AAGACATATT	TCTCAAGATC	AGAAGTACTA	TTCCAGTATG	2520
	GACGATTCAA	GGCTTGCTTC	ATAAACCAAG	GCAAGTAATA	GAGATTGGAG	TCTCTAAAAA	2580
10	GGTAGTTCCT	ACTGAATCTA	AGGCCATGCA	TGGAGTCTAA	GATTCAAATC	GAGGATCTAA	2640
	CAGAACTCGC	CGTGAAGACT	GGCGAACAGT	TCATACAGAG	TCTTTTACGA	CTCAATGACA	2700
	AGAAGAAAAT	CTTCGTCAAC	ATGGTGGAGC	ACGACACTCT	GGTCTACTCC	AAAAATGTCA	2760
15	AAGATACAGT	CTCAGAAGAC	CAAAGGGCTA	TTGAGACTTT	TCAACAAAGG	ATAATTTTCGG	2820
	GAAACCTCCT	CGGATTCCAT	TGCCCAGCTA	TCTGTCACTT	CATCGAAAGG	ACAGTAGAAA	2880
20	AGGAAGGTGG	CTCCTACAAA	TGCCATCATT	GCGATAAAGG	AAAGGCTATC	ATTCAAGATG	2940
	CCTCTGCCGA	CAGTGGTCCC	AAAGATGGAC	CCCCACCCAC	GAGGAGCATC	GTGGAAAAAG	3000
	AAGACGTTCC	AACCACGTCT	TCAAAGCAAG	TGGATTGATG	TGACATCTCC	ACTGACGTAA	3060
25	GGGATGACGC	ACAATCCCAC	TATCCTTCGC	AAGACCCTTC	CTCTATATAA	GGAAGTTCAT	3120
	TTCATTTGGA	GAGGACACGC	TGAAATCACC	AGTCTCTCTC	TATAAATCTA	TCTCTCTCTC	3180
30	TATAACCATG	GACCCAGAAC	GACGCCCCGGC	CGACATCCGC	CGTGCCACCG	AGGCGGACAT	3240
	GCCGGCGGTC	TGCACCATCG	TCAACCACTA	CATCGAGACA	AGCACGGTCA	ACTTCCGTAC	3300
	CGAGCCGCAG	GAACCGCAGG	AGTGGACGGA	CGACCTCGTC	CGTCTGCGGG	AGCGCTATCC	3360
35	CTGGCTCGTC	GCCGAGGTGG	ACGGCGAGGT	CGCCGGCATC	GCCTACGCGG	GGCCCTGGAA	3420
	GGCACGCAAC	GCCTACGACT	GGACGGCCGA	GTCGACCGTG	TACGTCTCCC	CCCGCCACCA	3480
40	GCGGACGGGA	CTGGGCTCCA	CGCTCTACAC	CCACCTGCTG	AAGTCCCTGG	AGGCACAGGG	3540
	CTTCAAGAGC	GTGGTCGCTG	TCATCGGGCT	GCCCAACGAC	CCGAGCGTGC	GCATGCACGA	3600
	GGCGCTCGGA	TATGCCCCCC	GCGGCATGCT	GCGGGCGGCC	GGCTTCAAGC	ACGGGAACTG	3660
45	GCATGACGTG	GGTTTCTGGC	AGCTGGACTT	CAGCCTGCCG	GTACCGCCCC	GTCCGGTCTT	3720
	GCCCCGTCACC	GAGATCTGAT	CTCACGCGTC	TAGGATCCGA	AGCAGATCGT	TCAAACATTT	3780
50	GGCAATAAAG	TTTCTTAAGA	TTGAATCCTG	TTGCCGGTCT	TGCGATGATT	ATCATATAAT	3840
	TTCTGTTGAA	TTACGTTAAG	CATGTAATAA	TTAACATGTA	ATGCATGACG	TTATTTATGA	3900
55	GATGGGTTTT	TATGATTAGA	GTCCCGCAAT	TATACATTTA	ATACGCGATA	GAAAACAAAA	3960

- 54 -

	TATAGCGCGC	AAACTAGGAT	AAATTATCGC	GCGCGGTGTC	ATCTATGTTA	CTAGATCGGG	4020
	AAGATCCTCT	AGAGTCGACC	TGCAGGCATG	CAAGCTTGGC	GTAATCATGG	TCATAGCTGT	4080
5	TTCCTGTGTG	AAATTGTTAT	CCGCTCACAA	TTCCACACAA	CATACGAGCC	GGAAGCATAA	4140
	AGTGTAAAGC	CTGGGGTGCC	TAATGAGTGA	GCTAACTCAC	ATTAATTGCG	TTGCGCTCAC	4200
	TGCCCCGCTTT	CCAGTCGGGA	AACCTGTCGT	GCCAGCTGCA	TTAATGAATC	GGCCAACGCG	4260
10	CGGGGAGAGG	CGGTTTGCGT	ATTGGGCGCT	CTTCCGCTTC	CTCGCTCACT	GACTCGCTGC	4320
	GCTCGGTTCGT	TCGGCTGCGG	CGAGCGGTAT	CAGCTCACTC	AAAGGCGGTA	ATACGGTTAT	4380
15	CCACAGAATC	AGGGGATAAC	GCAGGAAAGA	ACATGTGAGC	AAAAGGCCAG	CAAAGGCCA	4440
	GGAACCGTAA	AAAGGCCGCG	TTGCTGGCGT	TTTTCCATAG	GCTCCGCCCC	CCTGACGAGC	4500
	ATCACAAAAA	TCGACGCTCA	AGTCAGAGGT	GGCGAAACCC	GACAGGACTA	TAAAGATACC	4560
20	AGGCGTTTCC	CCCTGGAAGC	TCCCTCGTGC	GCTCTCCTGT	TCCGACCCTG	CCGCTTACCG	4620
	GATACCTGTC	CGCCTTTCTC	CCTTCGGGAA	GCGTGGCGCT	TTCTCAATGC	TCACGCTGTA	4680
25	GGTATCTCAG	TTCGGTGTAG	GTCGTTTCGT	CCAAGCTGGG	CTGTGTGCAC	GAACCCCCCG	4740
	TTCAGCCCGA	CCGCTGCGCC	TTATCCGGTA	ACTATCGTCT	TGAGTCCAAC	CCGGTAAGAC	4800
	ACGACTTATC	GCCACTGGCA	GCAGCCACTG	GTAACAGGAT	TAGCAGAGCG	AGGTATGTAG	4860
30	GCGGTGCTAC	AGAGTTCTTG	AAGTGGTGGC	CTAACTACGG	CTACACTAGA	AGGACAGTAT	4920
	TTGGTATCTG	CGCTCTGCTG	AAGCCAGTTA	CCTTCGGAAA	AAGAGTTGGT	AGCTCTTGAT	4980
35	CCGGCAAACA	AACCACCGCT	GGTAGCGGTG	GTTTTTTTGT	TTGCAAGCAG	CAGATTACGC	5040
	GCAGAAAAAA	AGGATCTCAA	GAAGATCCTT	TGATCTTTTC	TACGGGGTCT	GACGCTCAGT	5100
	GGAACGAAAA	CTCACGTTAA	GGGATTTTGG	TCATGAGATT	ATCAAAAAGG	ATCTTCACCT	5160
40	AGATCCTTTT	AAATTAAAAA	TGAAGTTTTA	AATCAATCTA	AAGTATATAT	GAGTAACTT	5220
	GGTCTGACAG	TTACCAATGC	TTAATCAGTG	AGGCACCTAT	CTCAGCGATC	TGTCTATTTT	5280
45	GTTTCATCCAT	AGTTGCCTGA	CTCCCCGTCG	TGTAGATAAC	TACGATACGG	GAGGGCTTAC	5340
	CATCTGGCCC	CAGTGCTGCA	ATGATACCGC	GAGACCCACG	CTCACC GGCT	CCAGATTTAT	5400
	CAGCAATAAA	CCAGCCAGCC	GGAAGGGCCG	AGCGCAGAAG	TGGTCCTGCA	ACTTTATCCG	5460
50	CCTCCATCCA	GTCTATTAAT	TGTTGCCGGG	AAGCTAGAGT	AAGTAGTTTCG	CCAGTTAATA	5520
	GTTTGCGCAA	CGTTGTTGCC	ATTGCTACAG	GCATCGTGGT	GTCACGCTCG	TCGTTTGGTA	5580
55	TGGCTTCATT	CAGCTCCGGT	TCCCAACGAT	CAAGGCGAGT	TACATGATCC	CCCATGTTGT	5640

- 55 -

5 GCAAAAAGC GGTTAGCTCC TTCGGTCCTC CGATCGTTGT CAGAAGTAAG TTGGCCGCAG 5700
TGTTATCACT CATGGTTATG GCAGCACTGC ATAATTCTCT TACTGTCATG CCATCCGTAA 5760
GATGCTTTTC TGTGACTGGT GAGTACTCAA CCAAGTCATT CTGAGAATAG TGTATGCGGC 5820
GACCGAGTTG CTCTTGCCCCG GCGTCAATAC GGGATAATAC CGCGCCACAT AGCAGAACTT 5880
10 TAAAAGTGCT CATCATTGGA AAACGTTCTT CGGGGCGAAA ACTCTCAAGG ATCTTACCGC 5940
TGTTGAGATC CAGTTCGATG TAACCCACTC GTGCACCCAA CTGATCTTCA GCATCTTTTA 6000
CTTTCACCAG CGTTTCTGGG TGAGCAAAAA CAGGAAGGCA AAATGCCGCA AAAAAGGGAA 6060
15 TAAGGGCGAC ACGGAAATGT TGAATACTCA TACTCTTCCT TTTTCAATAT TATTGAAGCA 6120
TTTATCAGGG TTATTGTCTC ATGAGCGGAT ACATATTTGA ATGTATTTAG AAAAATAAAC 6180
20 AAATAGGGGT TCCGCGCACA TTTCCCCGAA AAGTGCCACC TGACGTCTAA GAAACCATTA 6240
TTATCATGAC ATTAACCTAT AAAAATAGGC GTATCACGAG GCCCTTTCGT C 6291

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5560 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: T-DNA of plasmid pTHW142

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION:1..25
- (D) OTHER INFORMATION:/label= RB

/note= "right border sequence of octopine TL-DNA from pTiB6S3"

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION:complement (84..296)
- (D) OTHER INFORMATION:/label= 3'g7

/note= "3' untranslated region containing the polyadenylation signal of gene 7 of Agrobacterium T-DNA"

(ix) FEATURE:

- 56 -

(A) NAME/KEY: -
(B) LOCATION:complement (318..869)
(D) OTHER INFORMATION:/label= bar
/note= "region coding for posphinotricin acetyl
5 transferase"

(ix) FEATURE:

(A) NAME/KEY: -
(B) LOCATION:complement (830..2760)
10 (D) OTHER INFORMATION:/label= PSSU
/note= "promoter region of Rubisco small subunit gene of
Arabidopsis thali..."

(ix) FEATURE:

15 (A) NAME/KEY: -
(B) LOCATION:complement (2765..3058)
(D) OTHER INFORMATION:/label= 3'35S
/note= "3' untranslated region of the CaMV 35S transcript
containing polyadenylation signals"

(ix) FEATURE:

20 (A) NAME/KEY: -
(B) LOCATION:complement (3059..5056)
25 (D) OTHER INFORMATION:/label= uidA
/note= "region coding for beta-glucoronidase"

(ix) FEATURE:

30 (A) NAME/KEY: -
(B) LOCATION:complement (4483..4671)
(D) OTHER INFORMATION:/label= IV2
/note= "region corresponding to the second intron of the
ST-LS1 gene"

(ix) FEATURE:

35 (A) NAME/KEY: -
(B) LOCATION:complement (5067..5502)
(D) OTHER INFORMATION:/label= P35S
/note= "35S promoter region of CaMV"

(ix) FEATURE:

40 (A) NAME/KEY: -
(B) LOCATION:5533..5560
(D) OTHER INFORMATION:/label= LB
/note= "left border sequence of octopine TL-DNA from
45 pTIB6S3"

(ix) FEATURE:

50 (A) NAME/KEY: -
(B) LOCATION:5058..5059
(D) OTHER INFORMATION:/note= "region with unknown
sequence (may contain up to 20 nucleotides)"

(ix) FEATURE:

55 (A) NAME/KEY: -

- 57 -

(B) LOCATION:5077..5078

(D) OTHER INFORMATION:/note= "region with unknown
sequence (may contain up to 20 nucleotides)"

(ix) FEATURE:

(A) NAME/KEY: -

(B) LOCATION:5476..5479

(D) OTHER INFORMATION:/note= "region with unknown
sequence (may contain up to 20 nucleotides)"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

AATTACAACG	GTATATATCC	TGCCAGTACT	CGGCCGTCGA	GTACATGGTC	GATAAGAAAA	60
GGCAATTTGT	AGATGTTAAT	TCCCATCTTG	AAAGAAATAT	AGTTTAAATA	TTTATTGATA	120
AAATAACAAG	TCAGGTATTA	TAGTCCAAGC	AAAAACATAA	ATTTATTGAT	GCAAGTTTAA	180
ATTCAGAAAT	ATTTCAATAA	CTGATTATAT	CAGCTGGTAC	ATTGCCGTAG	ATGAAAGACT	240
GAGTGCAGATA	TTATGTGTAA	TACATAAATT	GATGATATAG	CTAGCTTAGC	TCATCGGGGG	300
ATCCTAGACG	CGTGAGATCA	GATCTCGGTG	ACGGGCAGGA	CCGGACGGGG	CGGTACCGGC	360
AGGCTGAAGT	CCAGCTGCCA	GAAACCCACG	TCATGCCAGT	TCCCGTGCTT	GAAGCCGGCC	420
GCCCGCAGCA	TGCCGCGGGG	GGCATATCCG	AGCGCCTCGT	GCATGCGCAC	GCTCGGGTCG	480
TTGGGCAGCC	CGATGACAGC	GACCACGCTC	TTGAAGCCCT	GTGCCTCCAG	GGACTTCAGC	540
AGGTGGGTGT	AGAGCGTGGA	GCCCAGTCCC	GTCCGCTGGT	GGCGGGGGGA	GACGTACACG	600
GTGCACTCGG	CCGTCCAGTC	GTAGGCGTTG	CGTGCCTTCC	AGGGGCCCCG	GTAGGCGATG	660
CCGGCGACCT	CGCCGTCCAC	CTCGGCGACG	AGCCAGGGAT	AGCGCTCCCG	CAGACGGACG	720
AGGTCGTCCG	TCCACTCCTG	CGGTCCTGTC	GGCTCGGTAC	GGAAGTTGAC	CGTGCTTGTC	780
TCGATGTAGT	GGTTGACGAT	GGTGCAGACC	GCCGGCATGT	CCGCCTCGGT	GGCACGGCGG	840
ATGTGGGCCG	GGCGTCGTTT	TGGGTCCATG	CAGTTAACTC	TTCCGCCGTT	GCTTGTGATG	900
GAAGTAATGT	CGTTGTTAGC	CTTGCGGGTG	GCTGGGAAGG	CAGCGGAGGA	CTTAAGTCCG	960
TTGAAAGGAG	CGACCATAGT	GGCCTGAGCC	GGAGAGGCAA	CCATAGTAGC	GGAAGAGAGC	1020
ATAGAGGAAG	CCATTGTTCT	TCTTTACTCT	TTGTGTGACT	GAGGTTTGGT	CTAGTGCTTT	1080
GGTCATCTAT	ATATAATGAT	AACAACAATG	AGAACAAGCT	TTGGAGTGAT	CGGAGGGTCT	1140
AGGATACATG	AGATTCAAGT	GGACTAGGAT	CTACACCGTT	GGATTTTGAG	TGTGGATATG	1200
TGTGAGGTTA	ATTTTACTTG	GTAACGGCCA	CAAAGGCCTA	AGGAGAGGTG	TTGAGACCCT	1260

- 58 -

	TATCGGCTTG	AACCGCTGGA	ATAATGCCAC	GTGGAAGATA	ATTCCATGAA	TCTTATCGTT	1320
	ATCTATGAGT	GAAATTGTGT	GATGGTGGAG	TGGTGCTTGC	TCATTTTACT	TGCCTGGTGG	1380
5	ACTTGGCCCT	TTCCTTATGG	GGAATTTATA	TTTTACTTAC	TATAGAGCTT	TCATACCTTT	1440
	TTTTTACCTT	GGATTTAGTT	AATATATAAT	GGTATGATTC	ATGAATAAAA	ATGGGAAATT	1500
10	TTTGAATTTG	TACTGCTAAA	TGCATAAGAT	TAGGTGAAAC	TGTGGAATAT	ATATTTTTTTT	1560
	CATTTAAAAG	CAAAATTTGC	CTTTTACTAG	AATTATAAAT	ATAGAAAAAT	ATATAACATT	1620
	CAAAATAAAA	TGAAAATAAG	AACCTTCAAA	AAACAGAACT	ATGTTTAATG	TGTAAAGATT	1680
15	AGTCGCACAT	CAAGTCATCT	GTTACAATAT	GTTACAACAA	GTCATAAGCC	CAACAAAGTT	1740
	AGCACGTCTA	AATAAACTAA	AGAGTCCACG	AAAATATTAC	AAATCATAAG	CCCAACAAAG	1800
20	TTATTGATCA	AAAAAAAAAA	ACGCCCAACA	AAGCTAAACA	AAGTCCAAAA	AAAACCTTCTC	1860
	AAGTCTCCAT	CTTCCTTTAT	GAACATTGAA	AACTATACAC	AAAACAAGTC	AGATAAATCT	1920
	CTTTCTGGGC	CTGTCTTCCC	AACCTCCTAC	ATCACTTCCC	TATCGGATTG	AATGTTTTTAC	1980
25	TTGTACCTTT	TCCGTTGCAA	TGATATTGAT	AGTATGTTTG	TGAAAATAAA	TAGGGTTAAC	2040
	AATCGAAGTC	ATGGAATATG	GATTTGGTCC	AAGATTTTCC	GAGAGCTTTC	TAGTAGAAAG	2100
30	CCCATCACCA	GAAATTTACT	AGTAAAATAA	ATCACCAATT	AGGTTTCTTA	TTATGTGCCA	2160
	AATTCAATAT	AATTATAGAG	GATATTTCAA	ATGAAAACGT	ATGAATGTTA	TTAGTAAATG	2220
	GTCAGGTAAG	ACATTAAAAA	AATCCTACGT	CAGATATTCA	ACTTTAAAAA	TTCGATCAGT	2280
35	GTGGAATTGT	ACAAAAATTT	GGGATCTACT	ATATATATAT	AATGCTTTAC	AACACTTGGA	2340
	TTTTTTTTTG	GAGGCTGGAA	TTTTTAATCT	ACATATTTGT	TTGGCCATG	CACCAACTCA	2400
40	TTGTTTAGTG	TAATACTTTG	ATTTTGTCAA	ATATATGTGT	TCGTGTATAT	TTGTATAAGA	2460
	ATTTCTTTGA	CCATATACAC	ACACACATAT	ATATATATAT	ATATATATTA	TATATCATGC	2520
	ACTTTTAATT	GAAAAAATAA	TATATATATA	TATAGTGCAT	TTTTTCTAAC	AACCATATAT	2580
45	GTTGCGATTG	ATCTGCAAAA	ATACTGCTAG	AGTAATGAAA	AATATAATCT	ATTGCTGAAA	2640
	TTATCTCAGA	TGTTAAGATT	TTCTTAAAGT	AAATTCTTTC	AAATTTTAGC	TAAAAGTCTT	2700
50	GTAATAACTA	AAGAATAATA	CACAATCTCG	ACCACGAAA	AAAAACACAT	AATAAATTTG	2760
	AATTAGCTTG	CATGCCTGCA	GGTCACTGGA	TTTTGGTTTT	AGGAATTAGA	AATTTTATTG	2820
	ATAGAAGTAT	TTTACAAATA	CAAATACATA	CTAAGGGTTT	CTTATATGCT	CAACACATGA	2880
55	GCGAAACCCT	ATAAGAACCC	TAATTCCTTT	ATCTGGGAAC	TACTCACACA	TTATTCTGGA	2940

- 59 -

GAAAAATAGA GAGAGATAGA TTTGTAGAGA GAGACTGGTG ATTTTGTGCGC CGGGTACCGA 3000
 GCTCGGTAGC AATCCCCGAG GCTGTAGCCG ACGATGGTGC GCCAGGAGAG TTGTTGATTC 3060
 ATTGTTTGCC TCCCTGCTGC GGTTTTTCAC CGAAGTTCAT GCCAGTCCAG CGTTTTTGGCA 3120
 GCAGAAAAGC CGCCGACTTC GGTTTGCGGT CGCGAGTGAA GATCCCTTTC TTGTTACCGC 3180
 CAACGCGCAA TATGCCTTGC GAGGTGCGAA AATCGGCGAA ATTCCATACC TGTTACCGA 3240
 CGACGGCGCT GACGCGATCA AAGACGCGGT GATACATATC CAGCCATGCA CACTGATACT 3300
 CTTCACTCCA CATGTCGGTG TACATTGAGT GCAGCCCGGC TAACGTATCC ACGCCGTATT 3360
 CGGTGATGAT AATCGGCTGA TGCAGTTTCT CCTGCCAGGC CAGAAGTTCT TTTCCAGTA 3420
 CCTTCTCTGC CGTTTCCAAA TCGCCGCTTT GGACATACCA TCCGTAATAA CGGTTACAGGC 3480
 ACAGCACATC AAAGAGATCG CTGATGGTAT CGGTGTGAGC GTCGCAGAAC ATTACATTGA 3540
 CGCAGGTGAT CGGACGCGTC GGGTCGAGTT TACGCGTTGC TTCCGCCAGT GGCGAAATAT 3600
 TCCCGTGAC TTGCGGACGG GTATCCGGTT CGTTGGCAAT ACTCCACATC ACCACGCTTG 3660
 GGTGGTTTTT GTCACGCGCT ATCAGCTCTT TAATCGCCTG TAAGTGCCTG TGCTGAGTTT 3720
 CCCCCTTGAC TGCCTCTTCG CTGTACAGTT CTTTCGGCTT GTTGCCCGCT TCGAAACCAA 3780
 TGCCTAAAGA GAGGTTAAAG CCGACAGCAG CAGTTTCATC AATCACCACG ATGCCATGTT 3840
 CATCTGCCCA GTCGAGCATC TCTTCAGCGT AAGGGTAATG CGAGGTACGG TAGGAGTTGG 3900
 CCCCATTCCA GTCCATTAAT GCGTGGTCGT GCACCATCAG CACGTTATCG AATCCTTTGC 3960
 CACGTAAGTC CGCATCTTCA TGACGACCAA AGCCAGTAA GTAGAACGGT TTGTGGTTAA 4020
 TCAGGAAGTG TTCGCCCTTC ACTGCCACTG ACCGGATGCC GACGCGAAGC GGGTAGATAT 4080
 CACACTCTGT CTGGCTTTTG GCTGTGACGC ACAGTTCATA GAGATAACCT TCACCCGGTT 4140
 GCCAGAGGTG CGGATTCACC ACTTGCAAAG TCCCGCTAGT GCCTTGTCCA GTTGCAACCA 4200
 CCTGTTGATC CGCATCACGC AGTTCAACGC TGACATCACC ATTGGCCACC ACCTGCCAGT 4260
 CAACAGACGC GTGGTTACAG TCTTGCGCGA CATGCGTCAC CACGGTGATA TCGTCCACCC 4320
 AGGTGTTCCG CGTGGTGTAG AGCATTACGC TGCGATGGAT TCCGGCATAG TTAAAGAAAT 4380
 CATGGAAGTA AGACTGCTTT TTCTTGCCGT TTTCGTCGGT AATCACCATT CCCGGCGGGA 4440
 TAGTCTGCCA GTTCAGTTCG TTGTTACAC AAACGGTGAT ACCTGCACAT CACCATGTTT 4500
 TGGTCATATA TTAGAAAAGT TATAAATTAA AATATACACA CTTATAAACT ACAGAAAAGC 4560

- 60 -

AATTGCTATA TACTACATTC TTTTATTTTG AAAAAAATAT TTGAAATATT ATATTACTAC 4620
TAATTAATGA TAATTATTAT ATATATATCA AAGGTAGAAG CAGAAACTTA CGTACACTTT 4680
5 TCCCGGCAAT AACATACGGC GTGACATCGG CTTCAAATGG CGTATAGCCG CCCTGATGCT 4740
CCATCACTTC CTGATTATTG ACCCACACTT TGCCGTAATG AGTGACCGCA TCGAAACGCA 4800
GCACGATACG CTGGCCTGCC CAACCTTTTCG GTATAAAGAC TTCGCGCTGA TACCAGACGT 4860
10 TGCCCGCATA ATTACGAATA TCTGCATCGG CGAACTGATC GTTAAACTG CCTGGCACAG 4920
CAATTGCCCCG GCTTTCTTGT AACGCGCTTT CCCACCAACG CTGATCAATT CCACAGTTTT 4980
15 CGCGATCCAG ACTGAATGCC CACAGGCCGT CGAGTTTTTT GATTTTCACGG GTTGGGGTTT 5040
CTACAGGACG GACCATGNNC CCGGGGATCC TCTAGANNTT ATAGAGAGAG AGATAGATTT 5100
ATAGAGAGAG ACTGGTGATT TCAGCGTGTC CTCTCCAAAT GAAATGAACT TCCTTATATA 5160
20 GAGGAAGGGT CTTGCGAAGG ATAGTGGGAT TGTGCGTCAT CCCTTACGTC AGTGGAGATG 5220
TCACATCAAT CCACTTGCTT TGAAGACGTG GTTGGAACGT CTTCTTTTTC CACGATGCTC 5280
25 CTCGTGGGTG GGGGTCCATC TTTGGGACCA CTGTCGGCAG AGGCATCTTG AATGATAGCC 5340
TTTCCTTTAT CGCAATGATG GCATTTGTAG GAGCCACCTT CCTTTTCTAC TGTCCTTTCG 5400
ATGAAGTGAC AGATAGCTGG GCAATGGAAT CCGAGGAGGT TTCCCGAAAT TATCCTTTGT 5460
30 TGAAAAGTCT CAATANNNG TCGACCTGCA GGCATGCAAG CTAATTCCGG GGAAGCTTAG 5520
ATCCATGGAG CCATTTACAA TTGAATATAT CCTGCCGCCG 5560

- 61 -

CLAIMS

1. A process for producing transgenic eucaryote cells which comprises:
contacting a culture of untransformed cells with an inhibitor of poly-
(ADP-ribose) polymerase, prior to transformation, for a period of time
sufficient to reduce the response of the cultured cells to stress and to
reduce the metabolism of said cultured cells, particularly to reduce the
electron flow in the mitochondrial electron transport chain; contacting
said untransformed cells with foreign DNA comprising at least one gene
of interest under conditions in which said foreign DNA is taken up by
said untransformed cells and said gene of interest is stably integrated in
the nuclear genome of said untransformed cells to produce said
transgenic cells; and
optionally recovering said transgenic cells from said culture.
2. The process of claim 1, wherein said eucaryotic cells are plant cells.
3. The process of claim 1 or 2, wherein said inhibitor is niacinamide,
preferably at a concentration of about 150 mg/l to 1000 mg/l, more preferably
at a concentration of about 200 mg/l to 500 mg/l, particularly at a concentration
of about 250 mg/l.
4. The process of any one of claims 1 to 3, wherein said untransformed cells
are cultured in a medium containing said inhibitor for a period of time of
approximately 2 to 28 days, preferably approximately 3 to 14 days, particularly
approximately 4 days prior to the contacting with said foreign DNA.
5. The process of any one of claims 1 to 4, wherein said cells contacted
with said foreign DNA are further cultured in a medium containing said inhibitor

- 62 -

for a period of time of approximately 1 to 14 days, preferably 2 to 4 days after contacting with said foreign DNA.

5 6. A process for increasing the frequency of obtaining transgenic plant cells which comprises:

contacting untransformed plant cells with foreign DNA comprising at least one gene of interest under conditions in which said foreign DNA is taken up by said untransformed cells and said gene of interest is stably integrated in the nuclear genome of said untransformed cells to produce
10 said transgenic cells

contacting cells with an inhibitor of poly-(ADP-ribose); and further culturing said cells in a medium containing said inhibitor for a period of time of approximately 1 to 14 days, preferably 1 to 4 days, particularly 1 day after contacting with said foreign DNA.

15 7. The process of any of claims 1 to 6, wherein said gene of interest comprises a promoter that directs expression selectively in certain cells or tissues of an eucaryotic organism.

20 8. The process of any one of claims 2 to 7, wherein said gene of interest comprises a promoter that directs expression selectively in stamen cells, particularly anther cells of a plant.

25 9. The process of claim 7 or 8, wherein said gene of interest encodes a protein that, when produced in a cell of an eucaryotic organism, kills or disables said cell.

10. The process of claim 9, wherein said gene of interest encodes a ribonuclease, particularly barnase.

- 63 -

11. The process of any one of claims 1 to 10, wherein a transgenic organism having said foreign DNA with said at least one gene of interest stably integrated in its genome is obtained from said transformed eucaryotic cell.

12. The process of claim 11, wherein said organism is a plant which is obtained by regeneration from a transformed plant cell.

13. The transgenic organism obtained by the process of claim 11 or 12.

14. A plant having foreign DNA integrated in the nuclear DNA of its cells only in the regions of said nuclear DNA that are transcriptionally active in said cells of said plant when said cells are treated with an effective amount of a PARP inhibitor for a period of time sufficient to reduce cell metabolism to a state where gene expression is essentially limited to genes expressed irrespective of the differentiated or physiological condition of the cell.

15. The plant according to claim 14, wherein said integration of the foreign DNA in said transcriptionally active region is verified by measuring the level of expressed mRNA corresponding to this foreign DNA when said cells are incubated in a medium containing a PARP-inhibitor.

16. The plant according to claim 14, wherein said transcriptionally active regions of the genome of said plant include regions which are minimally affected by cell differentiation or cell physiological and biochemical changes caused by external factors such as environmental conditions, especially stress conditions.

- 64 -

17. The plant or plant cell according to any one of the preceeding claims, wherein said plant or plant cell is a monocotyledonous plant or plant cell.

18. The plant or plant cell according to claim 17, wherein said plant or plant
5 cell is a cereal plant or plant cell.

19. The plant of plant cell according to claim 18, wherein said plant or plant cell is wheat or a wheat cell.

10 20. The plant according to any one of the preceeding claims, wherein said foreign DNA comprises a DNA sequence expressed selectively in specific tissues of said plant.

15 21. The plant of claim 20, wherein said foreign DNA comprises a DNA sequence encoding a cytotoxic molecule.

22. The plant of claim 21, wherein said foreign DNA comprises a DNA sequence encoding barnase.

20 23. A eucaryotic cell having foreign DNA integrated in its nuclear DNA only in the regions of said nuclear DNA that are transcriptionally active in said cell when said cell is treated with an effective amount of a PARP inhibitor for a period of time sufficient to reduce cell metabolism to a state where gene expression is essentially limited to genes expressed irrespective of the
25 differentiated or physiological condition of the cell.

COMBINED DECLARATION AND POWER OF ATTORNEY
FOR PATENT AND DESIGN APPLICATIONSATTORNEY DOCKET NO.
2121-127PPLEASE NOTE:
YOU MUST
COMPLETE THE
FOLLOWING:

As a below named inventor, I hereby declare that: my residence, post office address and citizenship are as stated next to my name; that I verily believe that I am the original, first and sole inventor (if only one inventor is named below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:*

Insert Title

GENETIC TRANSFORMATION USING A PARP INHIBITOR

Check Box If
Appropriate -
For Use Without
Specification
Attached

the specification of which is attached hereto unless the following box is checked:

☐ was filed on _____ as United States Application Number _____ or PCT International Application Number PCT/EP96/03366 filed July 31, 1996 and was amended on _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I do not know and do not believe the same was ever known or used in the United States of America before my or our invention thereof, or patented or described in any printed publication in any country before my or our invention thereof, or more than one year prior to this application, that the same was not in public use or on sale in the United States of America more than one year prior to this application, that the invention has not been patented or made the subject of an inventor's certificate issued before the date of this application in any country foreign to the United States of America on an application filed by me or my legal representatives or assigns more than twelve months (six months for designs) prior to this application, and that no application for patent or inventor's certificate on this invention has been filed in any country foreign to the United States of America prior to this application by me or my legal representatives or assigns, except as follows.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 (a)-(d) of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s)	Priority Claimed
95401844.6 United Kingdom 08/04/95	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
(Number) (Country) (Month/Day/Year Filed)	<input type="checkbox"/> Yes <input type="checkbox"/> No
(Number) (Country) (Month/Day/Year Filed)	<input type="checkbox"/> Yes <input type="checkbox"/> No
(Number) (Country) (Month/Day/Year Filed)	<input type="checkbox"/> Yes <input type="checkbox"/> No
(Number) (Country) (Month/Day/Year Filed)	<input type="checkbox"/> Yes <input type="checkbox"/> No
(Number) (Country) (Month/Day/Year Filed)	<input type="checkbox"/> Yes <input type="checkbox"/> No

I hereby claim the benefit under Title 35, United States Code, § 119(e) of any United States provisional application(s) listed below.

(Application Number)	(Filing Date)
(Application Number)	(Filing Date)

All Foreign Applications, if any, for any Patent or Inventor's Certificate Filed More Than 12 Months (6 Months for Designs) Prior To The Filing Date of This Application:

Country	Application No.	Date of Filing (Month/Day/Year)

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

(Application Number)	(Filing Date)	(Status — patented, pending, abandoned)
(Application Number)	(Filing Date)	(Status — patented, pending, abandoned)

I hereby appoint the following attorneys to prosecute this application and/or an international application based on this application and to transact all business in the Patent and Trademark Office connected therewith and in connection with the resulting patent based on instructions received from the entity who first sent the application papers to the attorneys identified below, unless the inventor(s) or assignee provides said attorneys with a written notice to the contrary:

RAYMOND C. STEWART (Reg. No. 21,066)
 JOSEPH A. KOLASCH (Reg. No. 22,463)
 JAMES M. SLATTERY (Reg. No. 28,380)
 DONALD C. KOLASCH (Reg. No. 23,038)
 CHARLES GORENSTEIN (Reg. No. 29,271)
 LEONARD R. SVENSSON (Reg. No. 30,330)
 MARC S. WEINER (Reg. No. 32,181)
 JOE McKINNEY MUNCY (Reg. No. 32,334)
 C. JOSEPH FARACI (Reg. No. 32,350)

TERRELL C. BIRCH (Reg. No. 19,382)
 ANTHONY L. BIRCH (Reg. No. 26,122)
 BERNARD L. SWEENEY (Reg. No. 24,448)
 MICHAEL K. MUTTER (Reg. No. 29,680)
 GERALD M. MURPHY, JR. (Reg. No. 28,977)
 TERRY L. CLARK (Reg. No. 32,644)
 ANDREW D. MEIKLE (Reg. No. 32,868)
 ANDREW F. REISH (Reg. No. 33,443)

PLEASE NOTE:
 YOU MUST
 COMPLETE THE
 FOLLOWING:

Send Correspondence to: **BIRCH, STEWART, KOLASCH AND BIRCH, LLP**

P.O. Box 747
 Falls Church, Virginia 22040-0747
 Telephone: (703) 205-8000
 Facsimile: (703) 205-8050

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full Name of First or Sole
 Inventor:
 Insert Name of Inventor
 Insert Date This
 Document Is Signed

Insert Residence
 Insert Citizenship

Insert Post Office
 Address

Full Name of Second
 Inventor, if any:

see above

Full Name of Third
 Inventor, if any:

see above

Full Name of Fourth
 Inventor, if any:

see above

Full Name of Fifth
 Inventor, if any:

see above

GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Marc	DE BLOCK		4/04/997
Residence (City, State & Country)		CITIZENSHIP	
Merelbeke, Belgium		Belgium	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
Abrikozenstraat 26, B-9820 Merelbeke, Belgium			
GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)		CITIZENSHIP	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)		CITIZENSHIP	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)		CITIZENSHIP	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)		CITIZENSHIP	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			

*Note: Must be completed
 — date this document is
 signed.

SEQUENCE LISTING

<110> DE BLOCK, MARC

<120> GENETIC TRANSFORMATION USING A PARP INHIBITOR

<130> 2121-0127P

<140> 08/817,188

<141> 1997-05-15

<150> PCT/EP96/03366

<151> 1996-07-31

<150> EP 95401844.6

<151> 1995-08-04

<160> 5

<170> PatentIn Ver. 2.0

<210> 1

<211> 4946

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: T-DNA of
plasmid pTHW107

<220>

<221> misc_feature

<222> Complement((1)..(25))

<223> T-DNA right border (RB)

<220>

<221> misc_feature

<222> Complement((97)..(330))

<223> 3'g7: 3' untranslated region containing the
polyadenylation signal of gene 7 of Agrobacterium
T-DNA

<220>

<221> misc_feature

<222> Complement((331)..(882))

<223> bar: region coding for phosphinotricin acetyl
transferase

<220>
 <221> misc_feature
 <222> Complement((883)..(2608))
 <223> promoter region of Rubisco small subunit gene of
 Arabidopsis thaliana (PSSU)

<220>
 <221> misc_feature
 <222> Complement((2658)..(3031))
 <223> 3' nos: 3' untranslated region containing the
 polyadenylation signal of the nopaline synthase
 gene of Agrobacterium T-DNA

<220>
 <221> misc_feature
 <222> Complement((3032)..(3367))
 <223> barnase: region coding for barnase

<220>
 <221> misc_feature
 <222> Complement((3368)..(4876))
 <223> PTA29: promoter region of TA29 gene of Nicotiana
 tabacum

<220>
 <221> misc_feature
 <222> Complement((4922)..(4946))
 <223> LB: T-DNA left border

<400> 1
 aattacaacg gtatatatcc tgccagtact cgcccgctga actcggccgt cgagtacatg 60
 gtcgataaga aaaggcaatt tgtagatgtt aattcccatc ttgaaagaaa tatagtttaa 120
 atatttatgt ataaaataac aagtcaggta ttatagtcca agcaaaaaca taaatttatt 180
 gatgcaagtt taaattcaga aatatttcaa taactgatta tatcagctgg tacattgccg 240
 tagatgaaaag actgagtgcg atattatgtg taatacataa attgatgata tagctagctt 300
 agctcatcgg gggatocctag acgcgtgaga tcagatctcg gtgacgggca ggaccggacg 360
 gggcggtacc ggcaggctga agtccagctg ccagaaaccc acgtcatgcc agttcccgtg 420
 cttgaagccg gccgcccga gcatgccgcg gggggcatat ccgagcgcct cgtgcatgcg 480
 cacgctcggg tcgttgggca gcccgatgac agcgaccacg ctcttgaagc cctgtgcctc 540
 cagggacttc agcaggtggg tgtagagcgt ggagcccagt cccgtccgct ggtggcgggg 600
 ggagacgtac acggtcgact cggccgtcca gtcgtaggcg ttgctgcctc tccaggggcc 660
 cgcgtaggcg atgccggcga cctcgccgtc cacctcggcg acgagccagg gatagcgctc 720
 ccgcagacgg acgaggtcgt ccgtccactc ctgcggttcc tgcggctcgg tacggaagtt 780
 gaccgtgctt gtctcgatgt agtggttgac gatggtgcag accgccggca tgtccgcctc 840
 ggtggcacgg cggatgtcgg cggggcgtcg ttctgggtcc attgttcttc tttactcttt 900
 gtgtgactga ggtttggtct agtgcttttg tcatctatat ataagtataa caacaatgag 960
 aacaagcttt ggagtgatcg gaggtctag gatacatgag attcaagtgg actaggatct 1020
 acaccgttgg attttgagtg tggatatgtg tgaggttaat tttacttggt aacggccaca 1080

aaggcctaag gagaggtggt gagaccctta tcggccttgaa ccgctggaat aatgccacgt 1140
ggaagataat tccatgaatc ttatcggtat ctatgagtga aattgtgtga tgggtggagt 1200
gtgcttgctc attttacttg cctgggtggac ttggcccttt ccttatggg aatttatatt 1260
ttacttacta tagagctttc ataccttttt ttaccttgg atttagttaa tatataatgg 1320
tatgattcat gaataaaaaat gggaaatttt tgaatttgta ctgctaaatg cataagatta 1380
ggtgaaactg tgggaatatat atttttttca tttaaaagca aaatttgcct tttactagaa 1440
ttataaatat agaaaaatat ataacattca aataaaaatg aaaataagaa ctttcaaaaa 1500
acagaactat gtttaatgtg taaagattag tcgcacatca agtcatctgt tacaatatgt 1560
tacaacaagt cataagccca acaaagttag cacgtctaaa taaactaaag agtccacgaa 1620
aatattacaa atcataagcc caacaaagt attgatcaaa aaaaaaaac gcccaacaaa 1680
gctaaacaaa gtccaaaaaa aacttctcaa gtctccatct tcctttatga acattgaaaa 1740
ctatacacia aacaagtcag ataaatctct ttctgggcct gtcttcccaa cctcctacat 1800
cacttcccta tcggattgaa tgttttactt gtaccttttc cgttgcaatg atattgatag 1860
tatgtttgtg aaaactaata gggtaacaa tcgaagtcatt ggaatatgga tttgggtccaa 1920
gattttccga gagctttcta gtagaaagcc catcaccaga aatttactag taaaataaat 1980
caccaattag gtttcttatt atgtgccaaa ttcaatataa ttatagagga tatttcaaat 2040
gaaaacgtat gaatgttatt agtaaaggt caggtaagac attaaaaaa tcctacgtca 2100
gatattcaac tttaaaaatt cgatcagtg ggaattgtac aaaaatttg gatctactat 2160
atatatataa tgctttacaa cacttggtatt tttttttgga ggctggaatt tttaatctac 2220
atatttgttt tggccatgca ccaactcatt gtttagtgta atactttgat tttgtcaaat 2280
atatgtgttc gtgtatattt gtataagaat ttctttgacc atatacacac acacatatat 2340
atatatatat atatattata tatcatgcac ttttaattga aaaaataata tatatatata 2400
tagtgcattt tttctaacia ccatatatgt tgcgattgat ctgcaaaaat actgctagag 2460
taatgaaaaa tataatctat tgctgaaatt atctcagatg ttaagatttt cttaaagtaa 2520
attctttcaa attttagcta aaagtcttgt aataactaaa gaataatata caatctcgac 2580
cacggaaaaa aaacacataa taaatttgaa tttcgaccgc ggtaccggga attcgagctc 2640
ggtaccgggg gatcttcccg atctagtaac atagatgaca ccgcgcgcga taatttatcc 2700
tagtttgccg gctatatttt gtttctatc gcgtattaaa tgtataattg cgggactcta 2760
atcataaaaa ccatctcat aaataacgtc atgcattaca tgtaattat tacatgctta 2820
acgtaattca acagaaatta tatgataatc atcgcaagac cggcaacagg attcaatctt 2880
aagaaacttt attgccaaat gtttgaacga tctgcttcgg atcctctaga gccggaaagt 2940
gaaattgacc gatcagagtt tgaagaaaaa tttattacac actttatgta aagctgaaaa 3000
aaacgcctc cgcaggaagc cgtttttttc gttatctgat ttttgtaaag gtctgataat 3060
ggtccgttgt tttgtaaate agccagtcgc ttgagtaaag aatccggtct gaatttctga 3120
agcctgatgt atagttaata tccgcttcac gccatgttcg tccgcttttg cccgggagtt 3180
tgccttccct gtttgagaag atgtctccgc cgatgctttt ccccgagcg acgtctgcaa 3240
ggttcccttt tgatgccacc cagccgaggg cttgtgcttc tgattttgta atgtaattat 3300
caggtagctt atgatatgtc tgaagataat ccgcaacccc gtcaaactg ttgataaccg 3360
gtaccatggg agctaatttc tttagtaaa aactttgatt tgagtgatga tgttgactg 3420
ttacacttgc accacaagg catatataga gcacaagaca tacacaacia cttgcaaaac 3480
taacttttgt tggagcattt cgaggaaaat ggggagtagc aggtcaatct gagggtaaca 3540
ttaaggtttc atgtattaat ttgttgcaaa catggactta gtgtgaggaa aaagtaccaa 3600
aattttgtct caccctgatt tcagttatgg aaattacatt atgaagctgt gctagagaag 3660
atgtttatc tagtccagcc acccacctta tgcaagtcg cttttagctt gattcaaaaa 3720
ctgatttaat ttacattgct aaatgtgcat acttcgagcc tatgtcgctt taattcgagt 3780
aggatgtata tattagtaca taaaaaatca tgtttgaatc atctttcata aagtgacaag 3840
tcaattgtcc cttcttgttt ggcactatat tcaatctgtt aatgcaaat atccagttat 3900
acttagctag atatccaatt ttgaataaaa atagctcttg attagtaaac cggatagtag 3960


```

caaagtcaca tatccatcaa acttctgggtg ctctgtggcta agttctgatc gacatgggggt 4020
taaaatttaa attgggacac ataaatagcc tattttgtgca aatctcccca tcgaaaatga 4080
cagattgtta catggaaaac aaaaagtcct ctgatagaag tcgcaaagta tcacaatttt 4140
ctatcgagag atagattgaa agaagtgcag ggaagcgggtt aactggaaca taacacaatg 4200
tctaaattaa ttgcattcgc taaccaaaaa gtgtattact ctctccggtc cacaataagt 4260
tatttttttg cccttttttt atggtccaaa ataagtgagt ttttttagatt tcaaaaatga 4320
tttaattatt tttttactac agtgcccttg gagtaaattgg tgttgagta tgtgttagaa 4380
atgtttatgt gaagaaatag taaagggttaa tatgatcaat ttcattgcta tttaattgta 4440
aaatgtgaat ttcttaatct gtgtgaaaac aaccaaaaaa tcacttattg tggaccggag 4500
aaagtatata aatatatatt tggaagcgac taaaaataaa cttttctcat attatacgaa 4560
cctaaaaaca gcatatggta gtttctaggg aatctaaatc actaaaatta ataaaagaag 4620
caacaagtat caatacatat gatttacacc gtcaaacacg aaattcgtaa atatttaata 4680
taataaagaa ttaatccaaa tagcctccca ccctataact taaactaaaa ataaccagcg 4740
aatgtatatt atatgcataa tttatatatt aaatgtgtat aatcatgtat aatcaatgta 4800
taatctatgt atatgggttag aaaaagtaaa caattaatat agccgggctat ttgtgtaaaa 4860
atccctaata taatcgcgac ggatccccgg gaattccggg gaagcttaga tccatggagc 4920
catttacaat tgaatatatc ctgccg                                     4946

```

<210> 2

<211> 6548

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: plasmid PTS172

<220>

<221> misc_feature

<222> Complement((2019)..(2288))

<223> 3' nos: 3' untranslated region containing the
polyadenylation signal of the nopaline synthase
gene of Agrobacterium T-DNA.

<220>

<221> misc_feature

<222> Complement((2289)..(2624))

<223> barnase: region coding for barnase

<220>

<221> misc_feature

<222> Complement((2625)..(4313))

<223> PE1: promoter region of E1 gene of rice

<220>

<221> misc_feature

<222> (4336)..(5170)

<223> P35S: 35S promoter region of Cauliflower Mosaic
Virus

<220>
 <221> misc_feature
 <222> Complement((5711)..(6262))
 <223> bar: region coding for phosphinotricin
 acetyltransferase

<220>
 <221> misc_feature
 <222> (6263)..(6496)
 <223> 3'g7: 3' untranslated region containing the
 polyadenylation signal of gene 7 of Agrobacterium
 T-DNA

<400> 2

aattcaagct	tgacgtcagg	tggcactttt	cggggaaatg	tgcgcggaac	ccctatttgt	60
ttatttttct	aaatacattc	aatatgtat	cgcctcatga	gacaataacc	ctgataaatg	120
cttcaataat	attgaaaaag	gaagagtatg	agtattcaac	atttccgtgt	cgcccttatt	180
cccttttttg	cggcattttg	ccttcctggt	tttgctcacc	cagaaacgct	ggtgaaagta	240
aaagatgctg	aagatcagtt	gggtgcacga	gtgggttaca	tcgaactgga	tctcaacagc	300
ggtaagatcc	ttgagagttt	tcgccccgaa	gaacgttttc	caatgatgag	cactttttaa	360
gtttctgctat	gtggcgcggt	attatcccgt	attgacgccg	ggcaagagca	actcggtcgc	420
cgcatacact	attctcagaa	tgacttggtt	gagtactcac	cagtcacaga	aaagcatctt	480
acggatggca	tgacagtaag	agaattatgc	agtgtctgca	taaccatgag	tgataacact	540
gcggccaact	tacttctgac	aacgatcgga	ggaccgaagg	agctaaccgc	ttttttgcac	600
aacatggggg	atcatgtaac	tcgccttgat	cgttggggaa	cggagctgaa	tgaagccata	660
ccaaacgacg	agcgtgacac	cacgatgcct	gtagcaatgg	caacaacggt	gcgcaaaacta	720
ttaactggcg	aactacttac	tctagcttcc	cggcaacaat	taatagactg	gatggaggcg	780
gataaagttg	caggaccact	tctgcgctcg	gcccttccgg	ctggctggtt	tattgctgat	840
aaatctggag	ccggtgagcg	tgggtctcgc	ggtatcattg	cagcactggg	gccagatggt	900
aagccctccc	gtatcgtagt	tatctacacg	acggggagtc	aggcaactat	ggatgaacga	960
aatagacaga	tcgctgagat	agggtgcctca	ctgattaagc	attggtaact	gtcagaccaa	1020
gtttactcat	atatacttta	gattgattta	aaacttcatt	tttaatttaa	aaggatctag	1080
gtgaagatcc	tttttggttc	gagtctcatg	acaaaaatcc	cttaacgtga	gttttcgttc	1140
cactgagcgt	cagaccccg	agaaaagatc	aaaggatctt	cttgagatcc	tttttttctg	1200
cgcgtaatct	gctgcttgca	aacaaaaaaa	ccaccgctac	cagcggtggt	ttgtttgccg	1260
gatcaagagc	taccaactct	ttttccgaag	gtaactggct	tcagcagagc	gcagatacca	1320
aatactgtcc	ttctagtgtg	gccgtagtta	ggccaccact	tcaagaactc	tgtagcaccg	1380
cctacatacc	tcgctctgct	aatcctgtta	ccagtggctg	ctgccagtgg	cgataagtcg	1440
tgtcttaccg	ggttggactc	aagacgatag	ttaccggata	aggcgacagc	gtcgggctga	1500
acgggggggt	cgtgcacaca	gccagcttg	gagcgaacga	cctacaccga	actgagatac	1560
ctacagcgtg	agcattgaga	aagcgccacg	cttcccgaag	ggagaaaggc	ggacaggtat	1620
ccggtgaagc	gcagggtcgg	aacaggagag	cgcacgaggg	agcttccagg	gggaaacgcc	1680
tggtatcttt	atagtcctgt	cgggtttcgc	cacctctgac	ttgagcgtcg	atttttgtga	1740
tgctcgtcag	gggggcggag	cctatggaaa	aacgccagca	acgcggcctt	tttacggttc	1800
ctggcctttt	gctggccttt	tgctcacatg	ttctttcctg	cgttatcccc	tgattctgtg	1860
gataaccgta	ttaccgcctt	tgagtgaagt	gataccgctc	gccgcagccg	aacgaccgag	1920
cgcagcgagt	cagtgaagca	ggaagcggaa	gagcgcccaa	tacgcaaacc	gcctctcccc	1980

gcgcgttggc ctgatcagaa ttcatatgca cgtgttcccg atctagtaac atagatgaca 2040
 ccgcgcgcga taatttatcc tagtttgccg gctatatattt gttttctatc gcgtattaaa 2100
 tgtataattg cgggactcta atcataaaaa cccatctcat aaataacgtc atgcattaca 2160
 tgttaattat tacatgctta acgtaattca acagaaatta tatgataatc atcgcaagac 2220
 cggcaacagg attcaatctt aagaaacttt attgccaaat gtttgaacga tctgcttcgg 2280
 aggttacctt atctgatttt tgtaaagggt tgataatggg ccgttgtttt gtaaatacagc 2340
 cagtcgcttg agtaaagaat ccggtctgaa tttctgaagc ctgatgtata gttaatatcc 2400
 gcttcacgcc atgttcgtcc gcttttgccc gggagtttgc cttccctgtt tgagaagatg 2460
 tctccgccga tgcttttccc cggagcgacg tctgcaagggt tcccttttga tgccaccag 2520
 ccgagggtt gtgcttctga ttttgtaatg taattatcag gtagcttatg atatgtctga 2580
 agataatccg caaccccgctc aaacgtgttg ataaccggta ccatcgcgac ggcttgatgg 2640
 atctcttgct ggacaccggg atgctaggat ggggttatcgt ggccggcgtg cgtgtgtggc 2700
 ttttgtaggc gccggcgacg gcgggggcaa tgtggcagggt gtagtcacggg gcaagcgtgc 2760
 gcaagtgaact gcaacaacca aggacggtca tggcgaaagc acctcacgcg tccaccgtct 2820
 acaggatgta gcagtagcac ggtgaaagaa gtgttgtccc gtccattagg tgcattctca 2880
 ccgttgggca gaacaggacc gttcaacagt taggttgagt gtaggacttt tacgtgggta 2940
 atgtatggca aatagtagta aattttgccc ccattgggtct ggctgagata gaacatatte 3000
 tggaaagcct ctagcatatc ttttttgaca gctaaacttt gcttcttgcc ttcttgggtct 3060
 agcaatgacg ttgcccattg cgtggcgaac atctggtaag gtaactgtat tctgttgttc 3120
 ccttcaacgg ctcaatcccc acaggccaag ctatcctttc cttggcagta taggtcctt 3180
 gagagattat actaccattt ttaagtgtct ataaagacga tgctctctaa ccagatcgat 3240
 cagaaacaca aagttttagc agcgtaatat cccacacaca tacacacacg aagctatgcc 3300
 tctcatttt ccgagagatt ctgacagtga ccagaatgtc agaatgccat ttcattgggca 3360
 caagtcgatc cacaagcttc ttgggtggagg tcaagggtgt ctattattat tctgtttcta 3420
 ggaaattatt cagaattagt gcctttttatc ataacttctc tctgagccga tgtggttttg 3480
 gatttcattg ttgggagcta tgcagttgcg gatattctgc tgtggaagaa caggaactta 3540
 tctgoggggg tccttgctgg ggcaacattg atatggttcc tgttcgatgt agtagaatac 3600
 aatataattc cgctcctttg ccagattgcc attcttgcca tgcttgatgt cttcatttgg 3660
 tcaaagccg caccactctt ggacaggat tagctttatt tcctgtggag atggtagaaa 3720
 actcagctta cagaaatggc atttcacgta gtataacgca agacattagg tactaaaact 3780
 caactaactg tttccgaatt tcagggtccc tccaaggatc ccagaaatca tcatctctga 3840
 acatgccttc agagaaatgg cattgaccgt ccattacaaa ctaacgtaca ctgtatctgt 3900
 tctttacgac attgcatgtg gaaaggatct gaagagattt ctcctggtac ataataatct 3960
 actcctttgc tacgttaata agagatgtaa aaacatgcaa cagttccagt gccaacattg 4020
 tccaaggatt gtgcaattct ttctggagcg ctaaaattga ccagattaga cgcacagaa 4080
 tattgaattg cagagttagc caataatcct cataatgtta atgtgctatt gttgttctact 4140
 actcaatata gttctggact aacaatcaga ttgtttatga tattaagggtg gttggatctc 4200
 tattggtatt gtcggcgatt ggaagttctt gcagcttgac aagtctacta tatattggta 4260
 ggtattccag ataaatatta aattttaata aaacaatcac acagaaggat ctgcccgcgc 4320
 tagcctaggc ccgggcccac aaaaatctga gcttaacagc acagttgctc ctctcagagc 4380
 agaatcgggt attcaacacc ctcatatcaa ctactacgtt gtgtataacg gtccacatgc 4440
 cggatatata gatgactggg gttgtacaaa ggccgcaaca aacggcgctt ccggagttgc 4500
 acacaagaaa tttgccacta ttacagaggc aagagcagca gctgacgct acacaacaag 4560
 tcagcaaca gacaggttga acttcatccc caaaggagaa gctcaactca agcccaagag 4620
 ctttgctaag gccctaaca gccacccaaa gcaaaaagcc cactgggtca cgctaggaac 4680
 caaaaggccc agcagtgatc cagcccaaaa agagatctcc tttgccccgg agattacaat 4740
 ggacgatttc ctctatcttt acgatctagg aagggaagttc gaagggtgaag gtgacgacac 4800
 tatgttcacc actgataatg agaagggttag cctcttcaat ttcagaaaga atgctgaccc 4860

```

acagatgggtt agagaggcct acgcagcagg tctcatcaag acgatctacc cgagtaacaa 4920
tctccaggag atcaaatacc ttccaagaa ggtaaagat gcagtcaaaa gattcaggac 4980
taattgcatc aagaacacag agaaagacat atttctcaag atcagaagta ctattccagt 5040
atggacgatt caaggcttgc ttcataaacc aaggcaagta atagagattg gagtctctaa 5100
aaaggtagtt cctactgaat ctaaggccat gcatggagtc taagattcaa atcgaggatc 5160
taacagaact cgccgtgaag actggcgaac agttcataca gagtctttta cgactcaatg 5220
acaagaagaa aatcttcgtc aacatgggtg agcacgacac tctgggtctac tccaaaaatg 5280
tcaaagatac agtctcagaa gaccaaaggg ctattgagac ttttcaacaa aggataat 5340
cgggaaacct cctcggattc cattgcccag ctatctgtca cttcatcgaa aggacagtag 5400
aaaaggaagg tggctcctac aaatgccatc attgcgataa aggaaaggct atcattcaag 5460
atgcctctgc cgacagtggc cccaaagatg gacccccacc cagcaggagc atcgtggaaa 5520
aagaagacgt tccaaccacg tcttcaaagc aagtggattg atgtgacatc tccactgacg 5580
taagggatga cgcacaatcc cactatcctt cgcaagaccc ttcctctata taaggaggtt 5640
catttcattt ggagaggaca cgctgaaatc accagtctct ctctataaat ctatctctct 5700
ctctataacc atggacccag aacgacgcc ggccgacatc cgccgtgcc aagaggcgga 5760
catgccggcg gtctgcacca tgcgtcaacca ctacatcgag acaagcacgg tcaacttccg 5820
taccgagccg caggaaccgc aggagtggac ggacgacctc gtccgtctgc gggagcgcta 5880
tccttggtc gtcgccgagg tggacggcga ggtcgccggc atcgccctac cgggcccctg 5940
gaaggcacgc aacgcctacg actggacggc cgagtcgacc gtgtacgtct ccccccgcga 6000
ccagcggacg ggactgggct ccacgctcta caccacactg ctgaagtccc tggaggcaca 6060
gggcttcaag agcgtgggtc gtgtcatcgg gctgcccac gacccgagcg tgcgcatgca 6120
cgaggcgctc ggatatgccc cccgcggcat gctgcgggcg gccggcttca agcacgggaa 6180
ctggcatgac gtgggtttct ggcagctgga cttcagcctg ccggtaccgc cccgtccggt 6240
cctgccgctc accgagatct gagatcacgc gttctaggat ccccgatga gctaagctag 6300
ctatatcatc aatttatgta ttacacataa tatcgactc agtctttcat ctacggcaat 6360
gtaccagctg atataatcag ttattgaaat atttctgaat ttaaacttgc atcaataaat 6420
ttatgttttt gcttggacta taatacctga cttgttattt tatcaataaa tatttaact 6480
atatttcttt caagatggga attaacatct acaaattgcc ttttcttacc gaccatgtac 6540
gtatcgcg
6548

```

<210> 3

<211> 1601

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: T72 promoter region

<220>

<221> misc_feature

<222> (1)..(1601)

<223> promoter region of T72 gene of rice

<400> 3

```

cgccgtgagt gtcttctgcc gccgaggggc tctcgctcgt cgctgatgcc tgcacgggtgc 60
gtgcgtgtgt gtcgtgggtg tgggtggcga acgcgacgcg agctcgattt ataggagggg 120
atcgaaggag gggagcgcgc gcggcgaggc ccgcgttgct cacctacgcc gcgcgcatgc 180

```

```

ggcggacgcg cggtcggcgc ccgcgccggc cgggaggacg agggcgcaag cgtgtgagcc 240
accgaacgcg cgcgcgcgcc gcggcgcgaa ctctccatcg cgtcgcggcg agccgagagc 300
cgacgagagc gtttcgcgcg cgcggttggg cggcgacaaa gatgggcccgt agccctgggc 360
ctcgtgccat cttttttttt cttttttgcc ttttttggc tggcaatttc tttttgtttt 420
tagtcttttt gtggtgataa tgtgtcgtct tccggtgaac taatttactc gttgatcttt 480
ttgtgtccct tcgaatatcc gcagtggtag aagatgacta ctactaccag tagttgatct 540
cgaatggcaa cttttgtgca gaacttattc cacggctatg tcagcttcca ctgtgactaa 600
aaaaactacg gccatctttt ggacttggtc tatcttggaa ctgaacaaaa aggacgatcc 660
tgatgtacac acggcatagt ttccagcact ggatgccaag ttgccaactg ttaccacgat 720
aatggaacga cgagatgaga tattatacaa gtccaatgga tcaagatcct gtgcagttgt 780
tattgttaact gtaacttaag ccgttaacat gtacatcaca tttcctactc tatcaatgtc 840
ttgtgcgggt tgtttcaaaa aaacatgtac atcacatgat ctagaacgga aggccaggat 900
atgaagtggg actgcagcaa aaacactgta gcagagatgt actattatgc atgtactgta 960
gcagtcactc agagccgttg gatctgaaaa cgaatggaca tgattgtgtg cagttgctat 1020
tgtgcagtta caatagcaac tgcatttgat cttaatccaa gtccaatata tgcagaacag 1080
tagctacgag ctggaaagga tgcaaatctg ggtgacactg acagcaaccg tggaagaaca 1140
acagcagcaa agtcccagag ggatggcaat ttgaaggaaat ttaaatactc taatattact 1200
ccaccggtta aaaaaaaciaa cttgctacgc ataatatatg ttccgattta tagcgagaag 1260
ttaatttttc atgagaagaa gaatatatat gtaatatgta ctaggagagt actcgcttca 1320
taaataataa tattcataag ttgtccagtg aagatagctt tagaaaaaac tagttatttt 1380
atgtgtcaaa ttttaaat tgaagtagtt agattatctt tctagtagtt ctgattgggt 1440
gaaaatgttt agattttcat gtgttaagag ttccgtatcc taaaaatagt aatataat 1500
taaatcatat atatatatat atatatatat atatatatat atatatatat 1560
tgttgaacgg tttgtgctct ggttgctatc ctgttctgtg g 1601

```

<210> 4

<211> 6291

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: plasmid pVE136

<220>

<221> misc_feature

<222> Complement((425)..(687))

<223> 3' nos: 3' untranslated region containing the
polyadenylation signal of the nopaline synthase
gene of Agrobacterium T-DNA

<220>

<221> misc_feature

<222> Complement((803)..(1138))

<223> barnase: region coding for barnase

<220>

<221> misc_feature

<222> Complement((1138)..(2317))

<223> PCa55: stamen-specific promoter from corn gene
CA55

<220>

<221> misc_feature

<222> (2355)..(3187)

<223> p35S: 35S promoter region of Cauliflower mosaic
virus

<220>

<221> misc_feature

<222> (3188)..(3739)

<223> bar: region coding for phosphinoacetyl transferase

<220>

<221> misc_feature

<222> (3757)..(4017)

<223> 3' nos: 3' untranslated region containing the
polyadenylation of the nopaline synthase gene of
Agrobacterium T-DNA

<220>

<221> misc_feature

<222> (699)..(702)

<223> region with unknown sequence (may contain up to 15
nucleotides)

<400> 4

```
tcgcgcggttt cggatgatgac ggtgaaaacc tctgacacat gcagctcccg gagacggtca 60
cagcttgtct gtaagcggat gccgggagca gacaagcccg tcagggcgcg tcagcgggtg 120
ttggcggggtg tcggggcttg cttactatg cggcatcaga gcagattgta ctgagagtgc 180
accatatgcg gtgtgaaata ccgcacagat gcgtaaggag aaaataccgc atcaggcgcc 240
attcgccatt caggctgcgc aactgttggg aaggcgcatc ggtgcggggc tcttcgctat 300
tacgccagct ggcgaaaggg ggatgtgctg caaggcgatt aagttgggta acgccagggt 360
tttcccagtc acgacgttgt aaaacgacgg ccagtgaatt cgagctcggg acccggggat 420
cttcccgatc tagtaacata gatgacaccg cgcgcgataa tttatcctag tttgcgcgct 480
atattttgtt ttctatcgcg tattaaatgt ataattgcgg gactctaata ataaaaaccc 540
atctcataaa taacgtcatg cattacatgt taattattac atgcttaacg taattcaaca 600
gaaattatat gataatcatc gcaagaccgg caacaggatt caatcttaag aaactttatt 660
gccaaatgtt tgaacgatct gcttcggatc ctctagagnn nnccggaaaag tgaaattgac 720
cgatcagagt ttgaagaaaa atttattaca cactttatgt aaagctgaaa aaaacggcct 780
ccgcaggaag ccgttttttt cgttatctga tttttgtaaa ggtctgataa tgggtccgtg 840
ttttgtaaat cagccagtcg cttgagtaaa gaatccgggtc tgaatttctg aagcctgatg 900
tatagttaat atccgcttca cgccatgttc gtccgctttt gcccgggagt ttgccttccc 960
tgtttgagaa gatgtctccg ccgatgcttt tccccggagc gacgtctgca aggttccctt 1020
ttgatgccac ccagccgagg gcttgtgctt ctgattttgt aatgtaatta tcaggtagct 1080
tatgatatgt ctgaagataa tccgcaaccc cgtcaaacgt gttgataacc ggtaccatgg 1140
ctgcagctag ttagctcgat gtatcttctg tatatgcagt gcagcttctg cgttttggct 1200
```

gctttgagct gtgaaatctc gctttccagt ccttgcgtgt tttatagtgc tgtacgttcg 1260
tgatcgtgag caaacagggc gtgcctcaac tactggtttg gttgggtgac aggcgccaac 1320
tacgtgctcg taaccgatcg agtgagcgta atgcaacatt ttttcttctt ctctcgatt 1380
ggtttcatcc agccaggaga cccgaatcga attgaaatca caaatctgag gtacagtatt 1440
tttacagtac cgttcgttcg aagggtcttcg acagggtcaag gtaacaaaat cagttttaaa 1500
ttgtttgttc agatcaaaga aaattgagat gatctgaagg acttgacct tcgtccaatg 1560
aaacacttgg actaattaga ggtgaattga aagcaagcag atgcaaccga aggtggtgaa 1620
agtggagttt cagcattgac gacgaaaacc ttcgaacggt ataaaaaaga agccgcaatt 1680
aaacgaagat ttgcaaaaa gatgcatcaa ccaagggag acgtgcatac atgtttgatg 1740
aaaactcgta aaaactgaag tacgattccc cattcccctc cttttctcgt ttcttttaac 1800
tgaagcaaag aatttgtatg tattccctcc attccatatt ctaggaggtt ttggcctttc 1860
ataccctcct ccatttcaaa ttatttgtca tacattgaag atatacacca ttctaattta 1920
tactaaatta cagcttttag atacatatat tttattatac acttagatac gtattatata 1980
aaacacctaa tttaaaataa aaaattatat aaaaagtgtg tctaaaaaat caaaatacga 2040
cataatttga aacggagggg tactacttat gcaaaccaat cgtggtaacc ctaaacccta 2100
tatgaatgag gccatgattg taatgcaccg tctgattaac caagatatca atggtcaaag 2160
atatacatga tacatccaag tcacagcgaa ggcaaatgtg acaacagttt tttttaccag 2220
agggacaagg gagaatatct attcagatgt caagttcccg tatcacactg ccaggtcctt 2280
actccagacc atcttccggc tctattgatg cataccagga attgatctag agtcgacctg 2340
caggcatgca agctcctacg cagcagggtc catcaagacg atctacccga gtaacaatct 2400
ccaggagatc aaataccttc ccaagaaggt taaagatgca gtcaaagat tcaggactaa 2460
ttgcatcaag aacacagaga aagacatatt tctcaagatc agaagtacta ttccagtatg 2520
gacgattcaa ggcttgcttc ataaaccaag gcaagtaata gagattggag tctctaaaaa 2580
ggtagttcct actgaatcta aggccatgca tggagtctaa gattcaaatc gaggatctaa 2640
cagaactcgc cgtgaagact ggcgaaacag tcatacagag tcttttacga ctcaatgaca 2700
agaagaaaat ctctgtcaac atggtggagc acgacactct ggtctactcc aaaaatgtca 2760
aagatacagt ctccagaagac caaagggtcta ttgagacttt tcaacaaagg ataatttcgg 2820
gaaacctcct cggattccat tgcccagcta tctgtcactt catcgaaagg acagtagaaa 2880
aggaaggtgg ctctacaaa tgccatcatt gcgataaagg aaaggctatc attcaagatg 2940
cctctgccga cagtgggtccc aaagatggac cccacccac gaggagcatc gtggaaaaag 3000
aagacgttcc aaccacgtct tcaaagcaag tggattgatg tgacatctcc actgacgtaa 3060
gggatgacgc acaatcccac tatccttcgc aagacccttc ctctatataa ggaagttcat 3120
ttcatttggg gaggacacgc tgaaatcacc agtctctctc tataaatcta tctctctctc 3180
tataaccatg gaccagaac gacgcccggc cgacatccgc cgtgccaccg aggcggacat 3240
gccggcggtc tgcaccatcg tcaaccacta catcgagaca agcacggtca acttccgtac 3300
cgagccgcag gaaccgcagg agtggacgga cgacctcgtc cgtctgcggg agcgctatcc 3360
ctggctcgtc gccgaggtgg acggcgaggt cgccggcatc gcctacgcgg gcccttgaa 3420
ggcacgcaac gcctacgact ggacggccga gtcgaccgtg tacgtctccc cccgccacca 3480
gcggacggga ctgggtcca cgctctacac ccacctgctg aagtccctgg aggcacaggg 3540
cttcaagagc gtggctcgtg tcatcgggct gcccacgac ccgagcgtgc gcatgcacga 3600
ggcgctcgga tatgcccccc gcggcatgct gcgggcggcc ggcttcaagc acgggaactg 3660
gcatgacgtg ggtttctggc agctggactt cagcctgccg gtaccgcccc gtccggtcct 3720
gcccgtcacc gagatctgat ctacgcgtc taggatccga agcagatcgt tcaaacattt 3780
ggcaataaag tttcttaaga ttgaatcctg ttgccgtct tgcgatgatt atcatataat 3840
ttctgttgaa ttacgttaag catgtaataa ttaacatgta atgcatgacg ttatttatga 3900
gatgggtttt tatgattaga gtcccgaat tatacattta atacgcgata gaaaacaaaa 3960
tatagcgcgc aaactaggat aaattatcgc gcgcgggtgc atctatgtta ctagatcggg 4020
aagatcctct agagtcgacc tgcaggcatg caagcttggc gtaatcatgg tcatagctgt 4080

```

ttcctgtgtg aaattgttat ccgctcacaa ttccacacaa catacgagcc ggaagcataa 4140
agtgtaaagc ctgggggtgcc taatgagtga gctaactcac attaattgcg ttgcgctcac 4200
tgcccgcgtt ccagtcggga aacctgtcgt gccagctgca ttaatgaatc ggccaacgcg 4260
cggggagagg cggtttgctg attgggcgct cttccgcttc ctcgctcact gactcgctgc 4320
gctcggtcgt tcggtgctg cgagcgggtat cagctcactc aaaggcggtat atacgggttat 4380
ccacagaatc aggggataac gcaggaaaga acatgtgagc aaaaggccag caaaaggcca 4440
ggaaccgtaa aaaggccgcg ttgctggcgt ttttccatag gtcgccccc cctgacgagc 4500
atcacaaaaa tcgacgctca agtcagaggt ggcgaaaccc gacaggacta taaagatacc 4560
aggcgtttcc ccctggaagc tccctcgtgc gctctcctgt tccgaccctg ccgcttaccg 4620
gatacctgtc cgcctttctc ccttcgggaa gcgtggcgct ttctcaatgc tcacgctgta 4680
ggtatctcag ttccgtgtag gtcgttcgct ccaagctggg ctgtgtgcac gaaccccccg 4740
ttcagcccga ccgctgcgcc ttatccggta actatcgtct tgagtccaac ccggttaagac 4800
acgacttatt gccactggca gcagccactg gtaacaggat tagcagagcg aggtatgtag 4860
gcggtgctac agagtctctg aagtgggtgc ctaactacgg ctacactaga aggacagtat 4920
ttggtatctg cgctctgctg aagccagtta ccttcggaaa aagagttggt agctcttgat 4980
ccggcaaaaa aaccaccgct ggtagcgggt gtttttttgt ttgcaagcag cagattacgc 5040
gcagaaaaaa aggatctcaa gaagatcctt tgatcttttc tacggggtct gacgctcagt 5100
ggaacgaaaa ctacggttaa gggattttgg tcatgagatt atcaaaaagg atcttcacct 5160
agatcctttt aaattaaaaa tgaagtttta aatcaatcta aagtatatat gagtaaaact 5220
ggtctgacag ttaccaatgc ttaatcagtg aggcacctat ctcagcgatc tgtctatttc 5280
gttcatccat agttgcctga ctcccgcgcg tgtagataac tacgatacgg gagggccttac 5340
catctggccc cagtgtctga atgataccgc gagaccacg ctcaccggct ccagatttat 5400
cagcaataaa ccagccagcc ggaagggccg agcgcagaag tggctcctgca actttatccg 5460
cctccatcca gtctattaat tgttgccggg aagctagagt aagtagttcg ccagttaata 5520
gtttgcgcaa cgttgttgcc attgctacag gcacgtgggt gtcacgctcg tcgtttggt 5580
tggcttcatt cagctccggt tcccaacgat caaggcgagt tacatgatcc cccatgttgt 5640
gcaaaaaagc ggtagctcc ttcggtcctc cgatcgttgt cagaagtaag ttggccgcag 5700
tgttatcact catggttatg gcagcactgc ataattctct tactgtcatg ccacccgtaa 5760
gatgcttttc tgtgactggt gactactcaa ccaagtcatt ctgagaatag tgtatgccc 5820
gaccgagttg ctcttgccc gcgtcaatac gggataatac cgcgccacat agcagaactt 5880
taaaagtgtc catcattgga aaacgttctt cggggcgaaa actctcaagg atcttaccgc 5940
tgttgagatc cagttcgatg taaccactc gtgcaccaa ctgatcttca gcatctttta 6000
ctttaccag cgtttctggg tgagcaaaaa caggaaggca aaatgccgca aaaaagggaa 6060
taagggcgac acggaaatgt tgaatactca tactcttct ttttcaatat tattgaagca 6120
tttatcaggg ttattgtctc atgagcggat acatatttga atgtatttag aaaaataaac 6180
aaataggggt tccgcgcaca tttccccgaa aagtgccacc tgacgtctaa gaaaccatta 6240
ttatcatgac attaacctat aaaaataggc gtatcacgag gccctttcgt c 6291

```

<210> 5

<211> 5560

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: T-DNA of
plasmid pTHW142

<220>

<221> misc_feature
 <222> (1)..(25)
 <223> RB: right border sequence of octopine TL-DNA from
 pTiB6S3

<220>
 <221> misc_feature
 <222> (84)..(296)
 <223> 3' g7: 3' untranslated region containing the
 polyadenylation signal of gene 7 of Agrobacterium
 T-DNA

<220>
 <221> misc_feature
 <222> (318)..(869)
 <223> bar: region coding for phosphinotricin
 acetyltransferase

<220>
 <221> misc_feature
 <222> (830)..(2760)
 <223> pSSU: promoter region of Rubisco small subunit
 gene of Arabidopsis thaliana

<220>
 <221> misc_feature
 <222> (2765)..(3058)
 <223> 3' untranslated region of the CaMV 35S transcript
 containing polyadenylation signals

<220>
 <221> misc_feature
 <222> (3059)..(5056)
 <223> uidA: region coding for beta-glucuronidase

<220>
 <221> misc_feature
 <222> (4483)..(4671)
 <223> IV2: region corresponding to the second intron of
 the ST-LS1 gene

<220>
 <221> misc_feature
 <222> (5067)..(5502)
 <223> P35S: 35S promoter region of CaMV

<220>
 <221> misc_feature

<222> (5533)..(5560)
 <223> LB: left border sequence of octopine TL-DNA from
 pTIB6S3

<220>
 <221> misc_feature
 <222> (5058)..(5059)
 <223> region with unknown sequence (may contain up to 20
 nucleotides)

<220>
 <221> misc_feature
 <222> (5077)..(5078)
 <223> region with unknown sequence (may contain up to 20
 nucleotides)

<220>
 <221> misc_feature
 <222> (5476)..(5479)
 <223> region with unknown sequence (may contain up to 20
 nucleotides)

<400> 5

```

aattacaacg gtatatatcc tgccagtact cggccgctcga gtacatggtc gataagaaaa 60
ggcaatttgt agatgttaat tcccatcttg aaagaaatat agtttaaata tttattgata 120
aaataacaag tcagggtatta tagtccaagc aaaaacataa atttattgat gcaagtttaa 180
attcagaaat atttcaataa ctgattatat cagctggtag attgccgtag atgaaagact 240
gagtgcgata ttatgtgtaa tacataaatt gatgatatag ctagcttagc tcatcggggg 300
atcctagacg cgtgagatca gatctcggtg acgggcagga ccggacgggg cggtaccggc 360
aggctgaagt ccagctgcca gaaaccacg tcatgccagt tcccgctgtt gaagccggcc 420
gcccgcagca tgccgcgggg ggcatatccg agcgcctcgt gcatgcgcac gctcgggtcg 480
ttgggcagcc cgatgacagc gaccacgctc ttgaagccct gtgcctccag ggacttcagc 540
aggtgggtgt agagcgtgga gccagctccc gtccgctggt ggcgggggga gacgtacacg 600
gtcgactcgg ccgtccagtc gtaggcgttg cgtgccttcc agggggccgc gtaggcgatg 660
ccggcgacct cgccgtccac ctccggcgacg agccagggat agcgctcccg cagacggacg 720
aggtcgctcg tccactcctg cggttcctgc ggctcggtag ggaagttgac cgtgcttgct 780
tcgatgtagt ggttgacgat ggtgcagacc gccggcatgt ccgcctcggg ggcacggcgg 840
atgtcggccg ggcgtcgttc tgggtccatg cagttaactc ttccgccgtt gcttgtgatg 900
gaagtaatgt cgttgtagc cttgcgggtg gctgggaagg cagcggagga cttaagtccg 960
ttgaaaggag cgaccatagt ggcctgagcc ggagaggcaa ccatagtagc ggaagagagc 1020
atagaggaag ccattgttct tctttactct ttgtgtgact gaggttttgt ctagtgcttt 1080
ggatcatctat atataatgat aacaacaatg agaacaagct ttggagtgat cggaggggtct 1140
aggatacatg agattcaagt ggactaggat ctacaccgtt ggatttttag tgtggatatg 1200
tgtgaggtta attttacttg gtaacggcca caaaggccta aggagaggtg ttgagaccct 1260
tatcggtctg aaccgctgga ataatgccac gtggaagata attccatgaa tcttatcggt 1320
atctatgagt gaaatttgtg gatgggtggg tgggtgcttg tcattttact tgcctggtgg 1380
acttggccct ttccttatgg ggaatttata ttttacttac tatagagctt tcataacctt 1440
tttttacctt ggatttagtt aatatataat ggtatgattc atgaataaaa atgggaaatt 1500

```

tttgaatttg	tactgctaaa	tgcataagat	taggtgaaac	tgtggaatat	atattttttt	1560
catttaaaag	caaaatttgc	cttttactag	aattataaat	atagaaaaat	atataacatt	1620
caaataaaaa	tgaaaataag	aacttttcaa	aaacagaact	atgttttaat	tgtaaagatt	1680
agtcgcacat	caagtcacat	gttacaatat	gttacaacaa	gtcataagcc	caacaaagtt	1740
agcacgtcta	aataaactaa	agagtccacg	aaaatattac	aatcataag	ccaacaaag	1800
ttattgatca	aaaaaaaaa	acgccaaca	aagctaaaca	aagtccaaaa	aaaacttctc	1860
aagtctccat	cttcctttat	gaacattgaa	aactatacac	aaaacaagtc	agataaatct	1920
ctttctgggc	ctgtcttccc	aacctcctac	atcacttccc	tatcggattg	aatgttttac	1980
ttgtaccttt	tccgttgcaa	tgatattgat	agtatgtttg	tgaaaactaa	taggggtaac	2040
aatcgaagtc	atggaatatg	gatttggtcc	aagattttcc	gagagctttc	tagtagaaag	2100
cccatcacca	gaaatttact	agtaaaataa	atcaccaatt	aggtttctta	ttatgtgcca	2160
aattcaatat	aattatagag	gatatttcaa	atgaaaacgt	atgaatgtta	ttagtaaag	2220
gtcaggtaag	acattaaaaa	aatcctacgt	cagatattca	actttaaaaa	ttcgatcagt	2280
gtggaattgt	acaaaaat	gggatctact	atatatatat	aatgctttac	aacacttgga	2340
tttttttttg	gaggctggaa	tttttaattc	acatatattgt	tttggccatg	caccaactca	2400
ttgttttagtg	taatactttg	attttgtcaa	atatatgtgt	tcgtgtatat	ttgtataaga	2460
atctctttga	ccatatacac	acacacatat	atatatatat	atatatatta	tatatcatgc	2520
acttttaatt	gaaaaataa	tatatatata	tatagtgc	tttttctaac	aaccatata	2580
gttgcgattg	atctgcaaaa	atactgctag	agtaatgaaa	aatataatct	attgctgaaa	2640
ttatctcaga	tgtaaagatt	ttcttaaagt	aaattctttc	aaatttttagc	taaaagtctt	2700
gtaataacta	aagaataata	cacaatctcg	accacggaaa	aaaaacacat	aataaatttg	2760
aattagcttg	catgcctgca	ggtcactgga	ttttggtttt	aggaattaga	aattttattg	2820
atagaagtat	tttacaataa	caaatacata	ctaagggttt	cttatatgct	caacacatga	2880
gcgaaaccct	ataagaacce	taattccctt	atctgggaac	tactcacaca	ttattctgga	2940
gaaaaataga	gagagataga	tttgtagaga	gagactggtg	atttttgcgc	cgggtaccga	3000
gctcggtagc	aattcccgcg	gctgtagccg	acgatggtgc	gccaggagag	ttgttgattc	3060
attgtttgcc	tccctgctgc	ggtttttcac	cgaagttcat	gccagtccag	cgtttttgca	3120
gcagaaaagc	cgccgacttc	ggtttgcggt	cgcgagtga	gatccctttc	ttgttaccgc	3180
caacgcgcaa	tatgccttgc	gaggtcgcaa	aatcggcgaa	attccatacc	tgttcaccga	3240
cgacggcgct	gacgcgatca	aagacgcggt	gatacatatc	cagccatgca	cactgatact	3300
cttcaactcca	catgtcgggtg	tacattgagt	gcagcccggc	taacgtatcc	acgccgtatt	3360
cgggtgatgat	aatcggctga	tgcatgtttc	cctgccaggc	cagaagttct	ttttccagta	3420
ccttctctgc	cgttttccaa	tcgccgcttt	ggacatacca	tccgtaataa	cggttcaggc	3480
acagcacatc	aaagagatcg	ctgatggtat	cgggtgtgagc	gtcgcagaac	attacattga	3540
cgcagggtgat	cggacgcgct	gggtcgagtt	tacgcgttgc	ttccgccagt	ggcgaatat	3600
tcccgtgcac	ttgcggacgg	gtatccggtt	cgttggcaat	actccacatc	accacgcttg	3660
ggtggttttt	gtcacgcgct	atcagctctt	taatcgcttg	taagtgcgct	tgctgagttt	3720
ccccgttgac	tgctctctcg	ctgtacagtt	ctttcggtt	gttgcccgt	tcgaaaccaa	3780
tgcttaaaga	gaggttaaag	cgcacagcag	cagtttcatc	aatcaccacg	atgccatggt	3840
catctgcccc	gtcagacatc	tcttcagcgt	aagggtaatg	cgaggtacgg	taggagttgg	3900
ccccaatcca	gtccattaat	gcgtggtcgt	gcacatcag	cacgttatcg	aatcctttgc	3960
cacgtaagtc	cgcattctca	tgacgaccaa	agccagtaaa	gtagaacggt	ttgtggttaa	4020
tcaggaactg	ttcgcccttc	actgccactg	accggatgcc	gacgcgaagc	gggtagatat	4080
cacactctgt	ctggcttttg	gctgtgacgc	acagttcata	gagataacct	tcacccgggt	4140
gccagagggtg	cggattcacc	acttgcaaag	tcccgcgtag	gccttgtcca	gttgcaacca	4200
cctgttgatc	cgcattcacgc	agttcaacgc	tgacatcacc	attggccacc	acctgccagt	4260
caacagacgc	gtggttacag	tcttgcgcg	catgcgtcac	cacggtgata	tcgtccacc	4320
aggtgttcgg	cgtggtgtag	agcattacgc	tgcatggat	tccggcatag	ttaaagaaat	4380

catggaagta	agactgcttt	ttcttgccgt	tttcgtcggt	aatcaccatt	cccgccggga	4440
tagtctgcca	gttcagttcg	ttgttcacac	aaacgggtgat	acctgcacat	caccatgttt	4500
tggatcatata	ttagaaaagt	tataaattaa	aatatacaca	cttataaaact	acagaaaagc	4560
aattgctata	tactacattc	ttttattttg	aaaaaaatat	ttgaaatatt	atattactac	4620
taattaatga	taattattat	atatatatca	aaggtagaag	cagaaactta	cgtaactttt	4680
tcccggcaat	aacatacggc	gtgacatcgg	cttcaaattg	cgtatagccg	ccctgatgct	4740
ccatcacttc	ctgattattg	acccacactt	tgccgtaatg	agtgaccgca	tcgaaacgca	4800
gcacgatacg	ctggcctgcc	caaccttttc	gtataaagac	ttcgcgctga	taccagacgt	4860
tgcccgcata	attacgaata	tctgcatcgg	cgaactgata	gttaaaactg	cctggcacag	4920
caattgcccg	gctttcttgt	aacgcgcttt	cccaccaacg	ctgatcaatt	ccacagtttt	4980
cgcgatccag	actgaatgcc	cacaggccgt	cgagtttttt	gatttcacgg	gttgggggtt	5040
ctacaggacg	gaccatgnnc	ccggggatcc	tctaganntt	atagagagag	agatagattt	5100
atagagagag	actggtgatt	tcagcgtgtc	ctctccaaat	gaaatgaact	tccttatata	5160
gaggaagggt	cttgccaagg	atagtgggat	tgtgcgtcat	cccttacgtc	agtggagatg	5220
tcacatcaat	ccacttgctt	tgaagacgtg	gttggaacgt	cttctttttc	cacgatgctc	5280
ctcgtgggtg	gggggtccatc	tttgggacca	ctgtcggcag	aggcatcttg	aatgatagcc	5340
tttcctttat	cgcaatgatg	gcatttgtag	gagccacctt	ccttttctac	tgtcctttcg	5400
atgaagtgac	agatagctgg	gcaatggaat	ccgaggaggt	ttcccgaat	tatcctttgt	5460
tgaaaagtct	caatannnng	tcgacctgca	ggcatgcaag	ctaattccgg	ggaagcttag	5520
atccatggag	ccattttacaa	ttgaatatat	cctgccgccg			5560